


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07D 295/18, A61K 31/445, C07C 259/06, A61K 31/16, 31/19, C07C 237/04, 237/06, C07D 233/32, 317/68, 213/74, 405/12</p>	A1	<p>(11) International Publication Number: WO 99/02510 (43) International Publication Date: 21 January 1999 (21.01.99)</p>
<p>(21) International Application Number: PCT/EP98/04220 (22) International Filing Date: 7 July 1998 (07.07.98) (30) Priority Data: 9714548.6 10 July 1997 (10.07.97) GB 9724395.0 18 November 1997 (18.11.97) GB (71) Applicant (for all designated States except US): PHARMACIA & UPJOHN S.P.A. [IT/IT]; Via Robert Koch, 1.2, I-20152 Milan (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): ALPEGIANI, Marco [IT/IT]; Via Tolmezzo, 12/5, I-20132 Milan (IT). BIS-SOLINO, Pierluigi [IT/IT]; Via Roma, 36/2, I-27020 San Giorgio di Lomellina (IT). ABRATE, Francesca [IT/IT]; Via Lodovico Settala, 45, I-20124 Milan (IT). PERRONE, Ettore [IT/IT]; Via Aldo Moro, 44, I-20010 Boffalora Ticino (IT). CORIGLI, Riccardo [IT/IT]; Via Morgantini, 31, I-20148 Milan (IT). JABES, Daniela [IT/IT]; Via Costanza, 3, I-20146 Milan (IT).</p>		<p>(81) Designated States: AL, AU, BR, CA, CN, CZ, HU, ID, IL, JP, KR, MX, NO, NZ, PL, RO, UA, US, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: MATRIX METALLOPROTEINASE INHIBITORS</p> <p>(57) Abstract</p> <p>A compound which is an amine derivative of formula (I) wherein W is -CONHOH or -COOH; R₁ and R₂ are each hydrogen or an organic residue, R₃ is an organic group, Q is a secondary or tertiary acyclic or cyclic amido group, and the pharmaceutically acceptable salts, solvates and hydrates thereof, are inhibitors of matrix metalloproteinases (MMPs) and of the release of tumor necrosis factor-α (TNF) from cells. They are therefore useful in the prevention, control and treatment of diseases in which MMPs or TNF are involved, especially tumoral and inflammatory diseases. Processes for their preparation and pharmaceutical compositions containing them are also described.</p> <div style="text-align: center;">  <p>(I)</p> </div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						


MATRIX METALLOPROTEINASE INHIBITORS

The present invention relates to new inhibitors of matrix metalloproteinases
5 (hereinafter MMPs), to a process for their preparation, to pharmaceutical compositions containing them, and to the use of such compounds in the prevention, control and treatment of diseases in which the proteolytic action of MMPs is involved.

Certain disease states are characterised by an imbalance of active MMPs and their natural inhibitors, the tissue inhibitors of metalloproteinases (hereinafter TIMPs). When TIMP levels
10 are insufficient, a progressive slow degradation of the extracellular matrix occurs, for example cartilage matrix loss in rheumatoid arthritis (L.A. Walakovits et al., *Arthritis Rheum.* 35:35-42, 1992) and osteoarthritis (D.D. Dean et al., *J. Clin. Invest.*, 84:678-685, 1989), and bone matrix degradation in osteoporosis (p.A. Hill et al., *Biochem. J.*, 308:167-175, 1995). In other situations, such as congestive heart failure, rapid degradation of the heart's
15 extracellular matrix occurs (P.W. Armstrong et al., *Canadian J. Cardiol.* 10:214-220, 1994). Cancer cells use MMPs, either expressed by themselves or by the surrounding tissues, to achieve rapid remodelling of the extracellular matrix. There is considerable evidence that MMPs are involved in at least 3 aspects of the growth and spread of tumors (e.g., see A.H. Davidson et al., *Chemistry & Industry*, 258-261, 1997, and references therein). In the process
20 of tumor metastasis, MMPs are used to break down the extracellular matrix, allowing primary tumor cancer cells to invade neighbouring blood vessels where they are transported to different organs and establish secondary tumors. The invasive growth at these secondary sites also needs MMPs to help break down tissue. In addition, MMP activity contributes to the invasive in-growth of new blood vessels (angiogenesis) which is required for tumors to
25 grow above a certain size.

Low molecular weight compounds able to inhibit one or more of the matrix metalloproteinases, in particular stromelysin-1 (MMP-3; EC 3.4.24.17), gelatinase A (MMP-2; EC 3.4.24.24), gelatinase B (MMP-9; EC 3.4.24.35), neutrophil collagenase or collagenase-2 (MMP-8; EC 3.4.24.34), interstitial collagenase or collagenase-1 (MMP-1; EC
30 3.4.27.7), matrilysin (MMP-7; EC 3.4.24.23), collagenase-3 (MMP-13), and the membrane-type metalloproteinase (MT-MMPs: MMP-14, MMP-15, MMP-16, MMP-17) are currently considered as promising therapeutic agents in degenerative, tumoral and autoimmune

pathologies (e.g., P.D. Brown: "Matrix metalloproteinase inhibitors: A new class of anticancer agent", *Curr. Opin. Invest. Drugs*, 2:617-626, 1993; A. Krantz: "Proteinases in Inflammation", *Annu. Rep. Med. Chem.* 28:187-195, 1993). Many of such compounds described hitherto are peptide derivatives or pseudopeptides, bearing analogies to recognized peptide substrates of these enzymes, and characterized in addition by a functional group capable of binding the Zn (II) atom present in the catalytic site of said enzymes. Known classes of MMP inhibitors include those in which the Zn binding group is a hydroxamic acid, and the skeleton, as represented in the general formula (A), mimicks the amino acid sequence of collagen at the site cleaved by collagenase:



wherein R_a , R_b , R_c , and R_d are hydrogen atoms or appropriate substituents (e.g., N.R.A. Beeley et al., "Inhibitors of matrix metalloproteinases (MMP's)", *Curr. Opin. Ther. Patents* 4:7-16, 1994; J.R. Porter et al., "Recent developments in matrix metalloproteinase inhibitors", *Exp. Opin. Ther. Patents* 5:1287-1296, 1995; J.R. Morphy et al., "Matrix metalloproteinase inhibitors: Current status", *Curr. Med. Chem.* 2:743-762, 1995; R.P. Beckett et al., "Recent advances in matrix metalloproteinase research", *DDT* 1:16-26, 1996). Said MMP inhibitors of the prior art can be described as "peptide-based hydroxamates" or "substrate-based" inhibitors (e.g., A.H. Davidson et al., "The inhibition of matrix metalloproteinase enzymes", *Chemistry & Industry*, 258-261, 1997).


Although MMPs have been recognized as drug targets for at least 20 years, and potent MMP inhibitors described by formula (A) have been disclosed since 1986 or before (e.g., see J.P. Dickens et al., U.S. Patent 4,599,361), no drug of this type has arrived at the market yet.

This is not because of questions about the therapeutic potential of MMP inhibitors, but because of problems of "peptide-based hydroxamates", such as aqueous solubility, metabolic stability, and other desirable properties, oral bioavailability in particular (e.g., J.R. Porter, reference above; J. Hodgson, "Remodelling MMPs", *Biotechnology* 13:554-557, 1995). For example, it is well known that most "peptide-based hydroxamates" of general formula (A)

are rapidly glucuronidated, oxidized to the carboxylic acid, and excreted in the bile (e.g., see J. Singh et al., Bioorg. Med. Chem. Lett. 5:337-342, 1995, and other references above). Finally, another type of problem of the known inhibitors described by general formula (A) may be one of tolerability. This problem is emerging for the most advanced MMP inhibitor in the clinic, marimastat (formula A; $R_a = OH$, $R_b = CH_2CHMe_2$, $R_c = CMe_3$, $R_d = Me$), which was reported to give musculoskeletal problems in humans. We have extended these observations by developing an animal model of tolerability with MMP inhibitors (S. Castellino et al., unpublished), involving intraperitoneal administration of the latter in rats for 10 consecutive days, and histological evaluation of stifle joints at the end of treatment. In this model, peptide-based MMP inhibitors of the prior art, e.g. Roche Ro31-9790 (formula A; $R_a = H$, $R_b = CH_2CHMe_2$, $R_c = CMe_3$, $R_d = Me$), at daily doses of 150 mg/kg or less, elicited hypertrophic fibrosis of stifle ligaments, interstitial hypertrophic fibrosis of skeletal muscles, hypertrophic fibroplasia of the periostium and synovium, and chondrosynplasia and decreased endochondral ossification of the epiphyseal plate. Although the precise reasons for these side-effects are not known at present, they support a strong need for better and diversified molecules, especially as far as the properties referred to above are concerned.

The present invention is concerned with novel MMP inhibitors, specifically characterised by the presence of a nitrogen atom as a substituent at the carbon atom next to the zinc-binding group, and with less or no peptidic character, as compared to substrate-based inhibitors of the prior art.

The present invention provides a compound which is an amine derivative of formula (I)



wherein

W is -CONHOH or -COOH;

R_1 and R_2 , which are the same or different, are each hydrogen or

- a group G, which is methyl, C₂-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₃-C₇ cycloalkyl, cycloalkyl-C₁-C₁₀- alkyl, aryl, aryl-C₁-C₁₀-alkyl, aryl-C₂-C₁₀-alkenyl, heterocyclyl, heterocyclyl-C₁-C₁₀-alkyl or heterocyclyl-C₂-C₁₀-alkenyl, the said methyl, alkyl, alkenyl, cycloalkyl, aryl and heterocyclyl groups being unsubstituted or substituted by one to three substituents; or
- 5 -SO₂-G, wherein G is as defined above; or
 -SO-G, wherein G is as defined above; or
 -CO-G, wherein G is as defined above; or
 -COO-G, wherein G is as defined above; or
 -SO₂-NH₂, -SO₂-NHG or -SO₂-NGG', wherein G is as defined above and G', which is the
 10 same or different, is as defined above for G, or G and G', together with the nitrogen atom to which they are attached, form a saturated or unsaturated 3- to 7-membered azaheterocyclic ring, which may be fused to a carbocyclic, heterocyclic, or aromatic ring, and may be substituted at any carbon or additional nitrogen atom, or
- a group -CONH₂, -CONHG or -CO-NGG' wherein G and G' are as defined above, or G
 15 and G', together with the nitrogen atom to which they are attached, constitute a saturated or unsaturated 3- to 7-membered azaheterocyclic ring, which may be fused to a carbocyclic, heterocyclic, or aromatic ring, and may be substituted at any carbon or additional nitrogen atom, or
- R₁ and R₂, taken together with the nitrogen atom to which they are attached, form a
 20 saturated or unsaturated 3- to 7-membered azaheterocyclic ring, which may be fused to a carbocyclic, heterocyclic, or aromatic ring, and may be substituted at any carbon or additional nitrogen atom;
- R₃ is C₁-C₁₅ alkyl, either unsubstituted or substituted by a C₃-C₇ cycloalkyl group, the alkyl and/or the cycloalkyl group being either unsubstituted, or substituted by one to three
 25 substituents selected from methyl, ethyl, C₃-C₄ linear or branched alkyl, fluoro, chloro, C₁-C₄ alkoxy, nitro, amino, dimethylamino, carboxy and carboxymethyl; or
- R₃ is a group -R-X-R^I wherein R is a chemical bond, -CH₂-, -(CH₂)_m- wherein m is an integer from 2 to 5, -CH=CH-, -CH₂CH=CH-, phenylene (i.e., -C₆H₄-), -CH₂CH=CH-C₆H₄-,
 -CH₂CH₂CH=CH-, -CH₂-CC-, -CH₂CH₂-CC-, -CH₂CH₂CH=CH-C₆H₄-, -CH₂-CC-C₆H₄-,
 30 or -CH₂CH₂-CC-C₆H₄-; X is a direct bond, an oxygen atom, a sulfur atom, or a sulfinyl -S(O)-, sulfonyl -S(O)₂ or carbamoyl group -CONH- or -NHCO-; and R^I is C₁-C₆ alkyl, C₂-C₆ alkenyl, phenyl, phenyl-(C₁-C₆)-alkyl, phenyl-(C₂-C₆)-alkenyl, heterocyclyl, or heterocyclyl-(C₁-C₆)-alkyl, either unsubstituted or substituted by a group selected from F, Cl,

Br, C₁-C₄ alkyl, C₁-C₄ alkoxy, alkylthio, arylthio, alkylsulfonyl, and arylsulfonyl;

Q, being a secondary or tertiary carboxamide, is:

- a group -CONHG or -CONGG', wherein G and G' are as defined above; or
- a group -CONH-CHGG', wherein G and G' are as defined above; or a group -CONG''-CHGG', wherein G'', being the same or different, is defined as G above; or
- a group -CONH-CH₂-CHGG' or a group -CONG''-CH₂-CHGG', wherein G, G' and G'' are as defined above; or
- a group -CO-azaheterocyclyl, wherein azaheterocyclyl, which is either unsubstituted or substituted, is as defined below; with the proviso that when Q is -CONHG, and G is methyl, alkyl-methyl, cycloalkyl-methyl, aryl-methyl or heterocyclyl-methyl, then such methyl or substituted methyl cannot be further substituted by a group -(CH₂)_t-CO₂H, wherein t is 0, or esters and amides thereof,

and a pharmaceutically acceptable salt, solvate or hydrate thereof.

As used herein the term "alkyl" refers to a straight or branched chain alkyl moiety having from 1 to 10 carbon atoms, including for example, methyl, ethyl, propyl, isopropyl, n-butyl, i-butyl, sec-butyl, tert-butyl, n-pentyl, isopentyl, n-hexyl, n-heptyl, n-octyl and so on.

The term "alkenyl" as used herein refers to a straight or branched chain alkenyl moiety having from 2 to 10 carbon atoms and having in addition one double bond of either E or Z stereochemistry where applicable. Examples of alkenyl groups are: vinyl, allyl, metallyl, butenyl, crotyl and so on.

The term "cycloalkyl" as used herein refers to a saturated carbocyclic group of 3 to 7 carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl.

The term "aryl" as used herein refers to a monocyclic or bicyclic aromatic hydrocarbon group of 6 to 10 carbon atoms, such as phenyl, naphthyl, indanyl; furthermore, "aryl" as used herein may refer to a diphenyl group (-C₆H₄-C₆H₅), a 4-pyridyl-phenyl group, and a methylenedioxyphenyl group.

The term "heterocyclyl" as used herein refers to a 3- to 7-membered, saturated or unsaturated heterocyclyl ring, containing at least one heteroatom selected from O, S and N, wherein any ring nitrogen may be oxidized as an N-oxide, any ring carbon may be oxidized as a carbonyl, and any ring sulfur may be oxidized as a sulfoxide or sulfone; and wherein said heterocyclyl ring may be optionally fused to a second 5- or 6-membered, saturated or unsaturated heterocyclyl ring, or to a C₃-C₇ cycloalkyl ring, or to a benzene or naphthalene ring. Examples of heterocyclyl groups are pyrrolyl, pyrrolidinyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, oxazolyl, isoxazolyl, thiazolyl,

isothiazolyl, thiadiazolyl, thienyl, tetrahydrothienyl, furyl, tetrahydrofuryl, aziridinyl, oxiranyl, azetidiny, succinimido, pyridyl, piperidinyl, pyrazinyl, piperazinyl, pyridazinyl, hexahydropyridazinyl, pyrimidinyl, pyranyl, tetrahydropyranyl, benzothienyl, benzothiazolyl, benzoxazolyl, isobenzofuranyl, benzofuranyl, benzimidazolyl, indazolyl, chromenyl, indolyl, oxindolyl, phthalimido, 1-oxo-2-isoindolyl, quinolyl, isoquinolyl, tetrahydroisoquinolyl, indoliziny, isoindolyl, 2-oxoisoindolyl, quinuclidinyl, hydantoinyl, saccharinyl, cinnolinyl, purinyl, morpholinyl, thiomorpholinyl, dioxanyl, dithianyl, azepinyl and so on.

The term "azaheterocyclyl" as used herein includes any of the heterocyclyl groups, as defined above, containing at least one nitrogen atom, said heterocyclyl group being linked to the rest of the molecule by a nitrogen atom.

Substituents which may be present in the methyl, alkyl, alkenyl, aryl, cycloalkyl, heterocyclyl and azaheterocyclyl groups in any of the above specifications include the following ones:

- a group $-(CH_2)_t-Hal$, wherein Hal is halo (i.e., fluoro, bromo, chloro or iodo), and t is an integer from 0 to 3;
- a group $-(CH_2)_t-CF_3$, or a group $-(CH_2)_t-CHF_2$, wherein t is as defined above;
- a group $-(CH_2)_t-OH$, wherein t is as defined above;
- a group $-(CH_2)_t-OR^I$, wherein t is as defined above, and R^I is straight or branched C_1-C_6 alkyl, aryl, arylmethyl, heterocyclyl, or heterocyclylmethyl, optionally substituted by hydroxy, methoxy, methyl, amino, methylamino, dimethylamino, chloro and fluoro;
- a group $-(CH_2)_t-OC(O)R^I$, wherein t and R^I are as defined above;
- a group $-(CH_2)_t-OC(O)OR^I$, wherein t and R^I are as defined above;
- a group $-(CH_2)_t-OC(O)NH_2$, or $-(CH_2)_t-OC(O)NHR^I$, or $-(CH_2)_t-OC(O)NR^IR^{II}$, wherein t is as defined above, R^I is as defined above, and R^{II} , being the same or different, is defined as R^I above; or R^I and R^{II} taken together with the nitrogen atom form an azaheterocycl ring;
- oxo;
- a group $-(CH_2)_t-NO_2$, wherein t is as defined above;
- a group $-(CH_2)_t-N_3$, wherein t is as defined above;
- a group $-(CH_2)_t-CN$, wherein t is as defined above;
- a group $-(CH_2)_t-SH$, wherein t is as defined above, and acetyl or phenylacetyl esters thereof (i.e., $-(CH_2)_t-SCOCH_3$ and $-(CH_2)_t-SCOCH_2C_6H_5$);
- a group $-(CH_2)_t-NH_2$, or $-(CH_2)_t-NHR^I$, or $-(CH_2)_t-NR^IR^{II}$, wherein t, R^I and R^{II} are as defined above, or R^I and R^{II} taken together with the nitrogen atom form an azaheterocyclyl ring;

- a group $-(CH_2)_t-NHC(O)R^{\text{II}}$, or $-(CH_2)_t-NR^{\text{II}}C(O)R^{\text{III}}$, or $-(CH_2)_t-NHC(O)OR^{\text{II}}$, wherein t, R^{II} and R^{III} are as defined above;
- a group $-(CH_2)_t-NH(CO)NH_2$, or $-(CH_2)_t-NH(CO)NHR^{\text{II}}$, or $-(CH_2)_t-NH(CO)NR^{\text{II}}R^{\text{III}}$, wherein t, R^{II} and R^{III} are as defined above, including the special case wherein R^{II} and R^{III} taken together with the nitrogen atom form an azaheterocyclyl ring;
- a group $-(CH_2)_t-NHSO_2R^{\text{II}}$, wherein t and R^{II} are as defined above;
- a group $-(CH_2)_t-NH(SO_2)NH_2$, or $-(CH_2)_t-NH(SO_2)NHR^{\text{II}}$, or $-(CH_2)_t-NH(SO_2)NR^{\text{II}}R^{\text{III}}$, wherein t, R^{II} and R^{III} are as defined above, including the special case wherein R^{II} and R^{III} taken together with the nitrogen atom form an azaheterocyclyl ring;
- a group $-(CH_2)_t-NHC(=NH)NH_2$, wherein t is as defined above;
- a group $-(CH_2)_t-CHO$, wherein t is as defined above;
- a group $-(CH_2)_t-C(O)R^{\text{II}}$, wherein t and R^{II} are as defined above;
- a group $-(CH_2)_t-CO_2H$, wherein t is as defined above, or esters or amides thereof, i.e., $-(CH_2)_t-CO_2R^{\text{II}}$, $-(CH_2)_t-CONH_2$, $-(CH_2)_t-CONHR^{\text{II}}$, $-(CH_2)_t-CONR^{\text{II}}R^{\text{III}}$, wherein R^{II} and R^{III} are as defined above, including the special case wherein R^{II} and R^{III} , taken together with the nitrogen atom form an azaheterocyclyl ring;
- a group $-(CH_2)_t-SO_3H$, wherein t is as defined above;
- a group $-(CH_2)_t-S(O)R^{\text{II}}$, wherein t and R^{II} are as defined above;
- a group $-(CH_2)_t-SO_2R^{\text{II}}$, wherein t and R^{II} are as defined above;
- a group $-(CH_2)_t-SO_2NH_2$, or $-(CH_2)_t-SO_2NHR^{\text{II}}$, or $-(CH_2)_t-SO_2NR^{\text{II}}R^{\text{III}}$, wherein t, R^{II} and R^{III} are as defined above;
- C_1-C_6 alkyl or C_2-C_6 alkenyl;
- C_3-C_7 cycloalkyl;
- phenyl, biphenyl (i.e., $-C_6H_4-C_6H_5$), methylenedioxyphenyl, methylenedioxyphenylmethyl (hereinafter piperonyl), benzyl, phenethyl, phenpropyl, naphthyl, naphthylmethyl, naphthylethyl, naphthylpropyl, either unsubstituted or substituted by one to three substituents selected from by C_1-C_4 alkyl, hydroxy, C_1-C_4 alkoxy, amino, methylamino, dimethylamino, chloro and fluoro.

When present carboxy, hydroxy, mercapto and amino groups may be either free or in a protected form. Protected forms of said groups are any of those generally known in the art, as described, for example, by T.W. Greene in "Protective Groups in Organic Chemistry", Wiley Interscience. Preferably, carboxy groups are protected as esters thereof, in particular methyl, ethyl, tert-butyl, benzyl, and 4-nitrobenzyl esters. Preferably, hydroxy groups are protected as ethers or esters thereof, in particular methoxymethyl ethers, tetrahydropyranyl ethers, benzyl ethers, acetates,

benzoates, pivalates. Preferably, mercapto groups are protected as thioethers or thioesters, in particular tert-butyl thioethers, thioacetates, thiobenzoates. Preferably, amino groups are protected as carbamates, e.g. tert-butoxycarbonyl and benzyloxycarbonyl derivatives, or as amides, e.g. acetamides and benzamides.

- 5 The present invention provides the salts of those compounds of formula (I) that have salt-forming groups, especially the salts of the compounds having an acid group, especially a carboxylic group, a N-hydroxycarbamoyl group, and a sulfo group, or the salts of the compounds having a basic group, especially an amino or guanidino group. The salts are especially physiologically tolerable salts, for example alkali metal and alkaline earth metal salts (e.g. sodium, potassium, lithium, calcium and magnesium salts), ammonium salts and salts with an appropriate organic amine or amino acid (e.g. arginine, procaine salts), and the addition salts formed with suitable inorganic acids (e.g. hydrochlorides, hydrobromides, sulfates, phosphates) or carboxylic and sulfonic organic acids (e.g. acetates, trifluoroacetates, citrates, succinates, malonates, lactates, tartrates, fumarates, maleates, methanesulfonates, *p*-toluenesulfonates). Some compounds of formula (I) which contain a carboxylate and an ammonium group may exist as zwitterions; such salts are also part of the present invention.
- 10


Furthermore, hydrates, solvates of compounds of formula (I), and physiologically hydrolyzable derivatives (i.e., prodrugs) of compounds of formula (I) are included within the scope of the present invention. Particularly preferred prodrugs of the compounds of formula (I) are ester derivatives. They include esters of compounds of formula (I) wherein W is -COOH, or wherein a carboxy group is present in any of the substituents R₁, R₂, R₃ and Q, which may be obtained by condensation of such carboxy group with a pharmaceutically acceptable alcohol, e.g. ethanol; or esters of compounds of formula (I) wherein a hydroxy group is present in any of the substituents R₁, R₂, R₃ and Q, which may be obtained by condensation of such hydroxy group with a pharmaceutically acceptable carboxylic acid, e.g. acetic acid, pivalic acid, benzoic acid and the like. Other particularly preferred prodrugs within the present invention are the cyclic condensation products between compounds of formula (I), wherein W is -CONHOH and R₁ is hydrogen, and formaldehyde, or an aldehyde of formula T-CHO, or a ketone of formula TT'CO, wherein T and T' are carbon radicals, such as lower alkyl, phenyl, benzyl, optionally substituted by one to three substituents selected from by C₁-C₄ alkyl, hydroxy, C₁-C₄ alkoxy, amino, methylamino, dimethylamino, chloro and fluoro. Such condensation products, which are represented herebelow, are obtained by mixing the two components, optionally in the presence of acid catalysts, as those employed for the formation of ketals from alcohols and ketones, and removing water by

20

25

30


evaporation, azeotropically or by molecular sieves.



- 5 The present invention encompasses all the possible stereoisomers (e.g. diastereoisomers, epimers, geometrical isomers) of the compounds of formula (I), as well as their racemic or optically active mixtures.

The present invention also includes, within its scope, pharmaceutical compositions comprising one or more of the compounds (I) as active ingredients, in association with pharmaceutically acceptable carriers, excipients or other additives, if desirable.

Preferred compounds within the present invention have the structure (I'):



- 15 wherein:

W is -CONHOH or -COOH;

R₁ and R₂ are:

- both hydrogen; or
- both C₁-C₄ alkyl, still preferably methyl; or

- 20 R₁ is hydrogen or methyl, and R₂ is a group G which is:

- C₁-C₁₀ alkyl or C₂-C₁₀ alkenyl, either unsubstituted or substituted by C₃-C₇ cycloalkyl, or by one to three substituents selected from chloro, fluoro, hydroxy, C₁-C₄ alkoxy, C₁-C₄ alkyl, amino, methylamino, dimethylamino, -CONH₂, -CONHCH₃ or -CONHC(CH₃)₃, C₁-C₄

- alkylthio, C₁-C₄ alkylsulfonyl, phenoxy, phenylthio and phenylsulfonyl, wherein the phenyl group may in turn be substituted by chloro, fluoro, methoxy or methyl; or
- C₃-C₇ cycloalkyl; or
 - an aryl group, more preferably phenyl, methylenedioxyphenyl, naphthyl or indanyl, each of which is optionally substituted by one to three substituents selected from chloro, fluoro, hydroxy, C₁-C₄ alkoxy, amino, methylamino, dimethylamino, C₁-C₄ alkyl, C₃-C₇ cycloalkyl, C₁-C₄ alkylthio, C₁-C₄ alkylsulfonyl, phenoxy, phenylthio and phenylsulfonyl, phenyl, benzyl, phenethyl, phenpropyl, naphthyl and pyridyl, and wherein any phenyl, naphthyl and pyridyl ring may in turn be substituted by one to three substituents selected from chloro, fluoro, methyl, hydroxy, methoxy, amino, methylamino and dimethylamino; or
 - an unsaturated heterocyclyl group selected from pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, thiadiazolyl, thienyl, furyl, pyridyl, pyrazinyl, pyridazinyl, pyrimidinyl, benzothienyl, benzothiazolyl, benzoxazolyl, isobenzofuranyl, benzofuranyl, benzimidazolyl, indazolyl, chromenyl, indolyl, oxindolyl, quinolyl, isoquinolyl, isoindolyl, cinnolyl and purinyl, each of which is optionally substituted by one to three substituents selected from chloro, fluoro, hydroxy, C₁-C₄ alkoxy, amino, methylamino, dimethylamino, C₁-C₄ alkyl, C₃-C₇ cycloalkyl, C₁-C₄ alkylthio, C₁-C₄ alkylsulfonyl, phenoxy, phenylthio and phenylsulfonyl, wherein the phenyl ring may be substituted by chloro, fluoro, methoxy or methyl; or
 - a saturated or partially saturated heterocyclyl group selected from pyrrolidinyl, tetrahydrothienyl, tetrahydrofuryl, aziridinyl, oxiranyl, azetidiny, piperidinyl, piperazinyl, hexahydropyridazinyl, tetrahydropyranyl, 1-oxo-2-isoindolyl, tetrahydroisoquinolyl, hydantoinyl, morpholinyl, thiomorpholinyl, dioxanyl, dithianyl, azepinyl and thiazolidinyl, wherein any ring nitrogen may be oxidized as an N-oxide, any ring carbon may be oxidized as a carbonyl, and any ring sulfur may be oxidized as a sulfoxide or sulfone; and the derivatives thereof substituted by one to three substituents selected from chloro, fluoro, hydroxy, C₁-C₄ alkoxy, amino, methylamino, dimethylamino, C₁-C₄ alkyl, C₃-C₇ cycloalkyl, C₁-C₄ alkylthio, C₁-C₄ alkylsulfonyl, phenoxy, phenylthio and phenylsulfonyl, wherein the phenyl ring may be substituted by chloro, fluoro, methoxy or methyl; or
 - C₁-C₁₀ alkyl, substituted by any of the unsaturated or saturated heterocyclyl groups as defined above, wherein any ring nitrogen may be oxidized as an N-oxide, any ring carbon may be oxidized as a carbonyl, and any ring sulfur may be oxidized as a sulfoxide or sulfone; or a derivative thereof substituted by one to three substituents selected from chloro, fluoro,

- hydroxy, C₁-C₄ alkoxy, amino, methylamino, dimethylamino, C₁-C₄ alkyl, C₃-C₇ cycloalkyl, -CONH₂, -CONHCH₃ and -CONHC(CH₃)₃, C₁-C₄ alkylthio, C₁-C₄ alkylsulfonyl, phenoxy, phenylthio and phenylsulfonyl, wherein the phenyl ring may be substituted by chloro, fluoro, methoxy or methyl; or
- 5 - C₁-C₁₀ alkyl, substituted by any of the aryl group as defined above, and the derivatives thereof substituted by one to three substituents selected from chloro, fluoro, hydroxy, C₁-C₄ alkoxy, amino, methylamino, dimethylamino, C₁-C₄ alkyl, C₃-C₇ cycloalkyl, C₁-C₄ alkylthio, C₁-C₄ alkylsulfonyl, phenoxy, phenylthio and phenylsulfonyl, wherein the phenyl ring may be substituted by chloro, fluoro, methoxy or methyl; or
- 10 R₁ is hydrogen or methyl, and R₂ is -SO₂-G, wherein G is as defined above; or
 R₁ is hydrogen or methyl, and R₂ is -CO-G, wherein G is as defined above; or
 R₁ is hydrogen or methyl, and R₂ is -CO-O-G, wherein G is as defined above; or
 R₁ is hydrogen or methyl, and R₂ is -SO₂-NH₂, -SO₂-NHG or -SO₂-NGG', wherein G is as defined above and G', which is the same or different, is as defined above for G; or
- 15 R₁ is hydrogen or methyl, and R₂ is -SO-NH-G, wherein G is as defined above; or
 R₁ is hydrogen or methyl, and R₂ is -CONHG or -CONGG', wherein G is as defined above and G', which is the same or different, is as defined above for G; or
 R₁ and R₂, taken together with the nitrogen atom to which they are attached, form a 3- to 7-membered azaheterocyclyl ring, optionally containing N, O, S or SO₂ as an additional ring
- 20 member, which may be substituted by oxo on one or two carbon ring atoms adjacent to the linking nitrogen atom, and which is optionally fused with a benzene ring, the azaheterocyclyl group being either unsubstituted or substituted at one or more carbon and/or nitrogen atoms by chloro, fluoro, hydroxy, C₁-C₄ alkoxy, amino, methylamino, dimethylamino, C₁-C₄ alkyl, phenyl, 4-fluorophenyl, benzyl, 4-fluorobenzyl, alpha-methylbenzyl, methylenedioxyphenyl, 2-phenethyl, 2-(4-fluorophenyl)ethyl, piperonyl, carbamoyl, -CONHCH₃, -CONHC(CH₃)₃, -CONH-(4-fluorophenyl), -CONH-pyridyl, -CONH-(methylenedioxy)phenyl -CONH-piperonyl;
- 25 or
 R₁ is hydrogen or methyl and R₂ is -SO₂-azaheterocyclyl, wherein azaheterocyclyl is as defined above; or
- 30 R₁ is hydrogen or methyl, and R₂ is -CO-azaheterocyclyl, wherein azaheterocyclyl is as defined above;
- R₃ is -CH₂-alkyl, -(CH₂)_n-cycloalkyl, -(CH₂)_n-O-alkyl, -(CH₂)_n-O-(CH₂)_m-cycloalkyl, -(CH₂)_n-O-(CH₂)_m-aryl, -(CH₂)_n-O-(CH₂)_m-heterocyclyl, -(CH₂)_n-S-alkyl, -(CH₂)_n-S-(CH₂)_m-

cycloalkyl, $-(CH_2)_n-S-(CH_2)_m$ -aryl, $-(CH_2)_n-S-(CH_2)_m$ -heterocyclyl, $-(CH_2)_n-SO$ -alkyl, $-(CH_2)_n-SO-(CH_2)_m$ -cycloalkyl, $-(CH_2)_n-SO-(CH_2)_m$ -aryl, $-(CH_2)_n-SO-(CH_2)_m$ -heterocyclyl, $-(CH_2)_n-SO_2$ -alkyl, $-(CH_2)_n-SO_2-(CH_2)_m$ -cycloalkyl, $-(CH_2)_n-SO_2-(CH_2)_m$ -aryl, $-(CH_2)_n-SO_2-(CH_2)_m$ -heterocyclyl, $-(CH_2)_n-CO$ -alkyl, $-(CH_2)_n-CO-(CH_2)_m$ -cycloalkyl, $-(CH_2)_n-CO-(CH_2)_m$ -aryl or $-(CH_2)_n-CO-(CH_2)_m$ -heterocyclyl, wherein alkyl, cycloalkyl, aryl and heterocyclyl are as defined above, and n and m, being the same or different, are zero or an integer of 1 to 5, and wherein the alkyl, cycloalkyl, aryl and heterocyclyl groups are optionally substituted by one to three substituents selected from chloro, fluoro, cyano, cyanomethyl, hydroxy, C₁-C₄ alkoxy, C₁-C₄ alkyl, C₁-C₄ alkylthio, C₁-C₄ alkylsulfonyl, phenyl, tolyl, 4-chlorophenyl, 4-fluorophenyl, 4-methoxyphenyl, 4-phenoxyphenyl, 4-(4-pyridyl)oxyphenyl, pyridyl; or R₃ is selected from isobutyl, n-hexyl, n-heptyl, n-octyl, n-nonyl and cyclopentylmethyl; or R₃ is selected from 2-phenylethyl, 3-phenylpropyl, 4-phenylbutyl, 5-phenylpentyl, wherein the phenyl group is either unsubstituted or substituted by chloro, fluoro, cyano, cyanomethyl, methyl, ethyl, propyl, butyl, mesyl, methoxy, ethoxy, propoxy, butoxy, phenoxy, 4-chlorophenoxy, 4-fluorophenoxy, benzyloxy, phenyl, 4-fluorophenyl, 4-chlorophenyl, 4-methoxyphenyl, 4-cyanophenyl, 4-cyanomethylphenyl, 4-pyridyl, 4-pyridyloxy; or R₃ is selected from phenylsulfonylmethyl or phenylsulphonylethyl, wherein the phenyl group is either unsubstituted or substituted by chloro, fluoro, cyano, cyanomethyl, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, butoxy, phenoxy, 4-chlorophenoxy, 4-fluorophenoxy, benzyloxy, phenyl, 4-fluorophenyl, 4-chlorophenyl, 4-methoxyphenyl, 4-cyanophenyl, 4-cyanomethylphenyl, 4-pyridyl, 4-pyridyloxy;

Q is:

- a group -CONHG, -CONGG', -CONH-CHGG', CON(CH₃)-CHGG', -CONH-CH₂-CHGG', or -CON(CH₃)-CH₂-CHGG', wherein G and G', being as defined above, are preferably selected from C₁-C₆ straight or branched alkyl, C₃-C₆ cycloalkyl, phenyl, tolyl, methylenedioxyphenyl, piperonyl and pyridyl, either unsubstituted or substituted by one to three substituents selected from chloro, fluoro, hydroxy, hydroxymethyl, C₁-C₄ alkoxy, amino, methylamino, dimethylamino, C₁-C₄ alkyl, C₃-C₇ cycloalkyl, -CONH₂, -CONHCH₃, -CONHC(CH₃)₃, -CONH(4-fluorophenyl), -CONH-pyridyl, -CONH-(methylenedioxy)phenyl, -CONH-piperonyl, carbomethoxy, carbethoxy, or a keto group -CO-R^{II}, wherein R^{II}, being as defined above, is selected from C₁-C₄ alkyl, phenyl, fluorophenyl, chlorophenyl, methylenedioxyphenyl, naphthyl, piperonyl, or a sulfone $-(CH_2)_n-SO_2-R^{II}$, wherein n and R^{II} are as defined above, or a sulfonamide $-(CH_2)_n-SO_2-NH_2$, $-(CH_2)_n-SO_2-NHR^{II}$, $-(CH_2)_n-SO_2-$

$\text{NR}^{\text{II}} \text{R}^{\text{III}}$, wherein n , R^{II} and R^{III} are as defined above, including the special case wherein R^{II} and R^{III} , taken together with the nitrogen atom to which they are attached, constitute an azaheterocyclyl ring, as defined above; or


- a group $-\text{CO}-\text{azaheterocyclyl}$, wherein azaheterocyclyl, being as defined above, is, either
 - 5 unsubstituted or substituted by one to three substituents selected from hydroxy, hydroxymethyl, C_1-C_4 alkoxy, carbamoyl, carbomethoxy, carbethoxy, mesyl, C_1-C_6 linear or branched alkyl, trifluoromethyl, C_3-C_7 cycloalkyl, aryl, heterocyclyl and aryl- $(\text{C}_1-\text{C}_3)\text{alkyl}$ or heterocyclyl- $(\text{C}_1-\text{C}_3)\text{alkyl}$; or said azaheterocyclyl group is substituted by a group $-\text{CONH}-\text{R}^{\text{II}}$, wherein R^{II} , being as defined above, is selected from methyl, isopropyl, tert-butyl,
 - 10 cyclopropyl, cyclopentyl, cyclohexyl, phenyl, 4-fluorophenyl, 4-chlorophenyl, 4-methoxyphenyl, 3,4-difluorophenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 4-piperidyl, 2-thiazolyl, 1-naphthyl, 2-naphthyl, 3-quinolyl, 5-quinolyl, 3-isoquinolyl, 5-isoquinolyl, 3-quinuclidinyl, methylenedioxyphenyl, piperonyl, 2-benzimidazolyl and 5-tetrazolyl; or said azaheterocyclyl group is substituted by a keto group $-\text{CO}-\text{R}^{\text{II}}$, or by a carbinol group of formula $-\text{CH}(\text{OH})-\text{R}^{\text{II}}$, wherein R^{II} , being as defined above, is selected from C_1-C_4 alkyl, phenyl, fluorophenyl, chlorophenyl, methylenedioxyphenyl, naphthyl, piperonyl, or by a sulfone $-(\text{CH}_2)_n-\text{SO}_2-\text{R}^{\text{II}}$, wherein n and R^{II} are as defined above, or by a sulfonamide $-(\text{CH}_2)_n-\text{SO}_2-\text{NH}_2$, $-(\text{CH}_2)_n-\text{SO}_2-\text{NHR}^{\text{II}}$, $-(\text{CH}_2)_n-\text{SO}_2-\text{NR}^{\text{II}} \text{R}^{\text{III}}$, wherein n , R^{II} and R^{III} are as defined above, including the special case wherein R^{II} and R^{III} , taken together with the nitrogen atom to which they are
 - 20 attached, constitute an azaheterocyclyl ring, as defined above;
- and the salts and solvates thereof.

When, in the above embodiment R_1 and R_2 form an azaheterocyclyl ring, the azaheterocyclyl ring is preferably selected from aziridine, azetidine, morpholine, thiomorpholine, piperidine, pyrrolidine, piperazine, thiazolidine, tetrahydroisoquinoline, hexahydropyridazine,

25 succinimido, phthalimido, saccharinyl, hydantoinyl, and oxoisoindolinyl. When R_2 is $-\text{CO}-\text{azaheterocyclyl}$ or $-\text{SO}_2-\text{azaheterocyclyl}$, the azaheterocyclyl moiety is preferably morpholino or piperidino. When Q is a group $-\text{CO}-\text{azaheterocyclyl}$, the azaheterocyclyl moiety is preferably selected from azetidine, morpholine, thiomorpholine, pyrrolidine, piperidine, piperazine, pyridazine, thiazolidine, tetrahydroisoquinoline, hexahydropyridazine and

30 hexamethylenimine. The aryl group or moiety in the definitions of Q is preferably selected from phenyl, 4-fluorophenyl, 4-methoxyphenyl, 4-methylphenyl, methylenedioxyphenyl, naphthyl, and heterocyclyl is preferably pyridyl.

A further preferred group of compounds are cyclic acetonide prodrugs of formula (II'):




wherein R₂, R₃ and Q are as defined above and T is methyl or a hydrogen atom, and T' is methyl, C₂-C₄ lower alkyl, phenyl, benzyl, optionally substituted by one to three substituents
 5 selected from by C₁-C₄ alkyl, hydroxy, C₁-C₄ alkoxy, amino, methylamino, dimethylamino, chloro and fluoro. More preferably, T and T' are methyl.

The present invention also provides a pharmaceutical composition comprising a pharmaceutically acceptable diluent or carrier and, as an active ingredient, a compound of the invention as defined above.

10 Compounds of the general formula (I) may be prepared by any suitable method known in the art, and/or by the following processes, which form another aspect of the invention. In the description and formulae below, the groups W, R₁, R₂, R₃ and Q are as defined above. It is understood that in the processes below any functional group (e.g. carboxyl, hydroxyl or amino), if needed or desired, can be masked by conventional methods and unmasked at the
 15 end or when convenient. Suitable protecting groups for such functionalities will be apparent to those skilled on the art and are well described in the chemical literature (see, for example: "Protective Groups in Organic Synthesis" by T.W. Greene, Wiley Interscience). It is also understood that any of the groups W, R₁, R₂, R₃ and Q can be converted by conventional methods into different groups W, R₁, R₂, R₃ and Q having any of the meaning previously
 20 defined, if desired, at the end or at any stage of the processes below. These conversions are known or will be apparent to those skilled in the art and are well described in the chemical literature (see, for example: "Comprehensive Organic Transformation" by R.C. Larock, VCH Publishers).

A preferred process for preparing a compound of formula (I) comprises:


25 (a) reacting a beta-lactam compound of formula (III):



wherein R_2 and R_3 are as defined above, and W' is COOH , CONHOH or a protected derivative thereof, with:

- a primary or secondary acyclic amine of formula G-NH_2 , GG'NH , GG'CH-NH_2 , GG'CH-NHCH_3 , $\text{GG'CH-CH}_2\text{-NH}_2$, or $\text{GG'CH-CH}_2\text{-NHCH}_3$, wherein G and G' are as defined above; or
- a cyclic saturated or unsaturated secondary amine, represented as azaheterocyclyl-H, wherein azaheterocyclyl is as defined above;

to obtain a compound of formula (IV):




wherein W' , R_2 , R_3 and Q are as defined above; and then:

- (b) converting said compound of formula (IV) into a compound of formula (I), wherein W , R_1 , R_2 , R_3 and Q are as defined above; and then:
- (c) if desired, removing the protecting groups and/or, if desired, converting any of the groups W , R_1 , R_2 , R_3 and Q into different groups W , R_1 , R_2 , R_3 and Q at the end or at any stage of the process.


It is evident that compounds with a desired configuration may be prepared starting from compounds (III) and (IV) with the appropriate configurations. Thus, a process for preparing

- preferred compounds of formula (I') comprises:

(a') reacting a beta-lactam compound of general formula (III'):




wherein R_2 and R_3 are as defined above, and W' is either COOH , CONHOH , or a protected derivative thereof, with an amine as defined above, to obtain a compound of formula (IV'):



5

wherein W' , R_2 , R_3 and Q are as defined above; and

(b') converting this compound of formula (IV') into a compound of formula (I'):




10 wherein W , R_1 , R_2 , R_3 and Q are as defined above.


The reaction between the beta-lactam of formula (III) or (III') and an amine among those hereabove detailed in step (a) or (a') above can be carried out in organic solvents, especially dimethylformamide (hereinafter DMF), tetrahydrofuran (hereinafter THF), acetonitrile, dimethylsulfoxide (hereinafter DMSO) and toluene, or in aqueous organic solvents, especially

15 aqueous THF, aqueous DMF, and aqueous acetonitrile, at temperatures ranging from 0 to 120 °C, either in the absence or in the presence of external bases. When the amine is a poor nucleophile, in order to accelerate the reaction, and achieving higher yields of the product of formula (IV) or (IV'), the reaction can be run in the presence of nucleophiles (NuH or salts thereof, wherein Nu is herebelow defined), which cleave the beta-lactam of formula (III) or

20 (III') more readily. In this case, an intermediate of formula (IIIa) or (III'a) is formed:



(IIIa)



(III'a)

wherein W' , R_2 and R_3 are as defined above, and Nu is selected from the group consisting of azido, imidazole, cyano, lower alkylthio, pyridylthio, phenylthio, and benzylthio. Said intermediate of formula (IIIa) or (III'a'), being an activated carboxylic acid derivative, reacts in the same milieu or in a separate step, and under the same reaction conditions, with the amine selected among those detailed above, giving rise to the product of formula (IV) or (IV'). Particularly preferred external nucleophiles are sodium azide, imidazole, and sodium and potassium cyanide; particularly preferred solvents are DMF and acetonitrile.

When in compounds of formula (III), (III'), (IV), (IV') above W' is a protected carboxy derivative, it is preferably benzyloxycarbonyl, p-nitrobenzyloxycarbonyl, p-methoxybenzyloxycarbonyl, tert-butoxycarbonyl, benzhydryloxycarbonyl, trityloxycarbonyl, trimethylsilyloxycarbonyl, tert-butyldimethyl-silyloxycarbonyl, phenyl-dimethyl-silyloxycarbonyl, allyloxycarbonyl, methoxycarbonyl, ethoxycarbonyl and n-butoxycarbonyl. When in compounds of formula (III), (III'), (IV), (IV') above W' is a protected derivative of CONHOH , it is preferably a group of formula CONHOR_{10} or $\text{CON(R}_{11}\text{)OR}_{10}$, wherein R_{10} and R_{11} are, respectively, hydroxy- and amino-protecting groups, known per se and removable by hydrogenolysis or by hydrolysis. Preferred R_{10} and R_{11} groups, which may be the same or different, include benzyl, p-methoxybenzyl, p-nitrobenzyl, trimethylsilyl, tert-butyl, tert-butoxycarbonyl, tetrahydropyranyl, and trityl.

The conversion of a compound of formula (IV) or (IV') into a compound of formula (I) or (I') in step (b) above may include any or all of the following steps in any order:

-(bⁱ): the conversion of the group W' , which is a protected carboxy or hydroxamate

derivative, into a group W, which is the unmasked carboxy or hydroxamic acid. This conversion is carried out by methodologies for unmasking of protective groups, which are well known in the art, as generally referred to above. A preferred conversion of this type is hydrogenolysis, especially in the presence of a palladium catalyst, in an inert organic solvent
5 such as ethanol or DMF or the like, especially at room temperature and under atmospheric pressure or moderate pressure, which is suitable for the conversion, e.g., of benzyl and p-nitrobenzyl esters into the parent carboxylic acids, or of O-benzyl and O,N-bis-benzyl hydroxamates into the parent hydroxamic acids. Another preferred conversion of this type is acid hydrolysis, especially by trifluoroacetic acid, hydrochloric acid, or by aluminium
10 trichloride, in the presence or absence of anisole, in inert organic solvents such as THF, acetonitrile and the like, especially between -20 and +30 °C, which is suitable for the conversion, e.g., of tert-butyl esters and p-methoxybenzyl esters into the parent carboxylic acids, or of O-tert-butylhydroxamates, O-(p-methoxybenzyl)-hydroxamates, and O,N-bis(p-methoxybenzyl) hydroxamates into the parent hydroxamic acids. A further preferred
15 conversion of this type is alkalyne hydrolysis, especially by NaOH, KOH, LiOH, $\text{KOSi}(\text{CH}_3)_3$, in an inert organic solvent or in water or in admixtures thereof, which is particularly suitable for the conversion of lower alkyl esters, e.g. the methyl, ethyl and n-butyl esters, into the parent carboxylic acids.

-(b''): the conversion of the group W', which is carboxy or an activated derivative thereof,
20 into a group W, which is -CONHOH. This conversion entails the condensation of such compounds of formula (IV) with hydroxylamine or a salt thereof, or with an O-protected hydroxylamine of formula $\text{R}_{10}\text{O-NH}_2$, or an N,O-diprotected hydroxylamine of formula $\text{R}_{10}\text{O-NHR}_{11}$, wherein R_{10} and R_{11} are as defined above, or a salt thereof, and then removal of said protecting groups R_{10} and R_{11} , if present, according to (b') above. Such condensation
25 is carried out according to general methodologies for the conversion of carboxylic acids or activated derivatives thereof into hydroxamic acids, which are well known in the art. In particular, activated derivatives of the carboxy group are the acid chloride, mixed anhydrides, and esters. In particular, the acid chloride is obtained by reacting the acid or a salt thereof with reagents such as oxalyl chloride or thionyl chloride; mixed anhydrides are obtained by
30 reacting the acid or a salt thereof with chlorocarbonates, such as ethyl chlorocarbonate, or with acid halides, such as pivaloyl chloride; esters, which are, preferably, the methyl, ethyl, n-butyl, pentafluorophenyl, hydroxysuccinyl, or hydroxybenzotriazolyl esters, are obtained by reaction of the acid with the corresponding alcohols in the presence of a dehydrating agent,

for example dicyclohexyl carbodiimide (hereinafter DCC), N,N-dimethylaminopropyl-N'-ethyl carbodiimide (EDC), and 2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline (EEDQ). Esters, in particular the methyl, ethyl, and n-butyl esters, may be present from the beginning in the azetidinone intermediates of formula (III) and (III') above. An O-protected hydroxylamine is, preferably, O-benzylhydroxylamine, O-tert-butylhydroxylamine, O-tert-butyl-diphenylsilyl hydroxyamine, O-(4-methoxybenzyl)-hydroxylamine, O-(4-nitrobenzyl)hydroxylamine, O-trimethylsilylhydroxylamine, and O-(tert-butoxycarbonyl)hydroxylamine. An N,O-diprotected hydroxylamine is, preferably, N,O-bis(benzyl)hydroxylamine, N,O-bis(4-methoxybenzyl)hydroxylamine, N,O-bis(tert-butoxycarbonyl)hydroxylamine, N-(tert-butoxycarbonyl)-O-(tert-butyl-dimethylsilyl)hydroxylamine, and N-(tert-butoxycarbonyl)-O-(tetrahydropyranyl)hydroxylamine. Preferably, the condensation reaction with hydroxylamine, O-protected hydroxylamines, N,O-diprotected hydroxylamines, and the salts thereof, is carried out in an inert organic solvent, such as DMF, THF, acetonitrile, dichloromethane, toluene and the like, at temperatures ranging from -20 to + 60 °C, optionally in the presence of a tertiary organic base, such as triethylamine and N-methylmorpholine. When protected hydroxylamines are employed, the protecting groups are removed after the condensation reaction, under the conditions described in (b') above;

-(bⁱⁱⁱ): the conversion of the group NHR_2 , being R_2 different from hydrogen, into a group NH_2 . This reaction can be carried out on compounds of formula (I), (I'), or on intermediates of formula (IV), (IV'), wherein R_2 is an amino protecting group, according to methods well known per se, for example by the methods of removal of amino protecting groups which are part of the techniques of peptide chemistry. Particularly preferred R_2 groups for such conversion are electron-withdrawing groups, in particular alkoxy- or benzyloxy-carbonyl groups such as tert-butoxycarbonyl, benzyloxycarbonyl and 4-nitro or 4-methoxy derivatives thereof, since the same particular R_1 groups efficiently assist the beta-lactam cleavage reaction detailed in step (a) above. In a preferred embodiment of the present invention, R_2 is tert-butoxycarbonyl, which can be removed by treatment with trifluoroacetic acid (TFA), optionally in the presence of anisole, in an inert organic solvent; in another preferred embodiment, R_2 is benzyloxycarbonyl or 4-nitrobenzyloxycarbonyl, which can be removed by catalytic hydrogenation;

-(b^{iv}): the conversion of the group NHR_2 , including the special case wherein R_2 is hydrogen, into a group NR_1R_2 , to be selected within the specifications stated above. Preferred R_1 and R_2 groups are the same groups detailed for the preferred compounds of


- formula (I). Such conversion encompasses functionalizations of amino groups well known in the art, such as alkylation, acylation, sulfonylation, and the like, and is performed according to methods well known per se. In a preferred embodiment of the present invention, such conversion is performed on compounds of formula (IV) or (IV') wherein W' is protected carboxy, thereafter removing the protecting group to obtain a compound of formula (I) or (I') wherein W is carboxy, by the general methodology described under (b') above and, optionally, by converting the so-obtained compound of formula (I) or (I') wherein W is carboxy into the corresponding compound wherein W is -CONHOH, by the general methodology described under (b'') above;
- 5 -(b^v): the conversion of any group W, R₁, R₂, R₃ and Q into any different group R₁, R₂, R₃ and Q, to be selected within the specifications stated above, by methodologies known per se. The resultant compounds of formula (I) or (I') may be converted into the desired salts, prodrugs, hydrates or solvates thereof by means of well known reactions, which include salt preparation by reaction with a pharmaceutically acceptable acid, or conversion of any hydroxy or carboxy group into an ester thereof, if desired, by condensation with a pharmaceutically acceptable alcohol or with a pharmaceutically acceptable carboxylic acid. In the particular case of the cyclic prodrugs described by formula (II) or (II'), said compounds are obtained from the corresponding compounds of formula (I) or (I') wherein W is -NHOH and R₁ is hydrogen, by reaction with an aldehyde of general formula T-CHO or a ketone of general formula TT'CO, wherein T and T' are as defined above, and removing water by evaporation.
- 10 The amines used in step (a) above are known compounds or are prepared from known compounds by known methods.
- 15 The beta-lactams of formula (III) or (III') above are known compounds or can be prepared from known compounds by methodologies known per se or by analogy with the specific preparative examples herein. In particular, a preferred preparation of compounds of formula (III) or (III') includes:
- 20 -(dⁱ): cyclization of an aspartic acid derivative to obtain a compound of formula (III) or (III') wherein R₃ is hydrogen, by reaction with a suitable condensing agent;
- 25 -(dⁱⁱ): conversion of a compound of formula (III) or (III') wherein R₃ is hydrogen into a compound of formula (III) or (III') wherein R₃ is as described above, by deprotonation with a strong base and alkylation of the resulting beta-lactam enolate with an agent of formula R₃-X, wherein X is halo, e.g. chloro, bromo or iodo, or sulfonyloxy, e.g. triflate, mesylate or the like.
- 30

General conditions for step (dⁱ) above are described in the literature, the preferred aspartic acid derivative being usually dibenzyl aspartate or di(4-nitro)benzyl aspartate. Some of the resultant azetidinones (III) or (III') wherein R₃ is hydrogen, W' is carboxy or an ester thereof, and R₂ is hydrogen or tert-butyldimethylsilyl are also commercially available. A preferred compound in step (dⁱⁱ) is a compound of formula (III) or (III') wherein R₃ is hydrogen, R₂ is tert-butyldimethylsilyl, and W' is free carboxy, since the carboxylate anion helps at avoiding racemization at the azetidinone 4-C, and increases regioselectivity of alkylation by the R₃-X reagent at the azetidinone 3-C.


The conditions described in steps (a), (b), (c) and (d) do not usually promote epimerization or racemization at pre-existing chiral centers. Thus, when the aspartic acid derivative used as a reagent in step (dⁱ) above is an L-aspartic acid derivative, the reaction affords a chiral azetidinone of formula (III'), wherein R₃ is hydrogen and R₂, W' are as described above. In such chiral azetidinone of formula (III') wherein R₃ is hydrogen, the configuration at the C-4 atom addresses stereoselective alkylation, by the R₃-X reagent, in a transoid relationship with the W' substituent. Therefore, starting from such intermediates of formula (III'), step (dⁱⁱ) above stereoselectively affords the corresponding azetidinones of formula (III') wherein R₃, being as described above, is different from hydrogen, with the depicted configurations at the two chiral centers. Said configurations of the two chiral centers are the same as found in compounds of formula (I') herein specifically preferred. Accordingly, it can be appreciated that steps (dⁱ) and (dⁱⁱ) above are essential part of an original, fully stereocontrolled route to the compounds of formula (I').

Another process for the preparation of compounds of formula (I) comprises:

(e) alkylation of an aspartic acid derivative of formula (V):



wherein W' is either carboxy or a protected derivative thereof, Q' is either Q as described above, or carboxy or protected carboxy, and R₁, R₂ are as described above, with a reagent of formula R₃-X, wherein R₃ and X are as described above, to obtain a compound of formula (VI):




wherein W' , R_1 , R_2 , R_3 and Q' are as described above; and:

(f) conversion of such compound of formula (VI) into the desired compound of formula (I),

5 wherein W , R_1 , R_2 , R_3 and Q are as described above.


We have observed that the configuration at the chiral carbon atom in compounds of formula (V) addresses the configuration at the new chiral center in the product of formula (VI) in an anti-fashion, exclusively or predominantly. Thus, starting from an L-aspartic derivative of formula (V'):

10



wherein W' , R_1 , R_2 , R_3 and Q' are as described above, step (e) predominantly or exclusively affords a compound of formula (VI'):

15



wherein W' , R_1 , R_2 , R_3 and Q' are as described above, from which the preferred compounds of formula (I') are obtained by step (f) above.

20 The reaction of step (e) above is carried out according to conventional carbanion chemistry, i.e. with 2-4 equivalents of a strong base, such as LDA and the like, in aprotic organic solvents, such as THF, DMF, N-methylpyrrolidone, HMPA and the like, at temperatures

ranging from -70°C to room temperature. A preferred W' group is free carboxy. A preferred R₁ group is hydrogen. A preferred R₂ group is tert-butoxycarbonyl. Preferred Q' groups are either the acyclic or cyclic tertiary carboxamides described within Q above, or a carboxy ester, which in step (f) can be converted into the group Q above as generally known for the preparation of carboxamides.

The compounds of formula (I) and (I') provided by the present invention are characterized by high inhibitory activity on matrix metalloproteinases (MMPs), especially collagenases, gelatinases and stromelysins. For example, the following protocol was used to assess the biochemical activity of compounds of formula (I') against MMP-1, MMP-2, and MMP-3 (respectively, human interstitial collagenase, gelatinase A, and stromelysin-1).

BIOCHEMICAL ASSAY (Protocol A)

Human collagenase (MMP-1) was obtained as truncated recombinant enzyme encompassing residues 101-269 and did not require activation. Human gelatinase-A (MMP-2) was obtained as pro-enzyme (72 kDa) and was activated with 1 mM 4-aminophenylmercuric acetate for 30 min at 37 °C immediately prior to use. Human stromelysin-1-255 (MMP-3) was obtained as a recombinant pro-enzyme isolated from *E. coli* and activated by heat (1 h, 55 °C).

The substrate was the commercial quenched-fluorescence peptide Mca-Pro-Leu-Gly-Leu-Dpa-Ala-Arg-NH₂, that is (7-methoxycoumarin-4-yl)Acetyl-Pro-Leu-Gly-Leu-(3-[2,4-dinitrophenyl]-L-2,3-diamino-propionyl)-Ala-Arg-NH₂ (C.G. Knight et al., FEBS Lett. 296:263, 1992). The enzymes cleave at the Gly-Leu bond, removing the internally quenching Dpa group. Release of the highly fluorescent peptide Mca-Pro-Leu was followed fluorimetrically at 37°C using a Perkin Elmer LS-50 Fluorescence Spectrophotometer fitted with a thermostatted four position stirring cell changer. The excitation wavelength was set at 326 nm (bandwidth 5 nm) and the emission at 392 nm (bandwidth 20 nm). K_m values of this substrate for the three MMP's is 70 μM or more (Knight et al., reference above). Substrate concentration was 2 μM in the tests, so that we could approximate to unit the term $(1 + [\text{substrate}] / K_M)$ in calculations. The substrate was stable for over 60 minutes in the assay conditions, giving no appreciable increment of fluorescence. Full response was adjusted against 200 nM Mca-Pro-Leu-OH (the released fluorescent peptide) and the instrument was calibrated in the range 0-100 nM Mca-Pro-Leu-OH, corresponding to 0-5% extent of hydrolysis of the 2 μM substrate. The aqueous assay buffer was 50 mM Tris/HCl pH=7.4 containing 0.15 M NaCl, 10 mM CaCl₂, 0.01 mM ZnCl₂ and 0.05% Brij 35. The concentration of the activated enzyme was determined by titration against representative

inhibitors described in the literature and synthesized in house: Roche Ro31-9790 (N.G. Knebel et al., J. Chromatogr. B 673:213, 1995) for MMP-1, and Celltech CT-1418 (compound 1 in S.K. Chander, J. Pharm. Sci. 84:404, 1995) for MMP-2 and MMP-3. Enzyme concentrations in the tests were set at 1.0 nM for MMP-1, 0.04 nM for MMP-2, and 3.0 nM for MMP-3. Under our assay conditions, we measured k_{cat} / K_M values of 26900, 669000 and 9740 $1/(M \times s)$ for MMP-1, MMP-2 and MMP-3, respectively. All the three enzymes were found stable for over three hours in the assay conditions.

The inhibition constant for compounds of the present invention was measured at steady state (K_i^* ; see J.F. Morrison and C.T. Walsh, Adv. Enzymol. Relat. Areas Mol. Biol. 61:201, 1988), upon preincubation experiment, by measuring V_0 , the initial rate in the absence of inhibitor, and V_s , the steady-state velocity, at different concentrations of inhibitors in the region of their enzyme-inhibitor dissociation constants.

On a routine basis, 1.94 mL of assay buffer was pre-heated at 37°C in a vial, 0.02 mL of inhibitor in DMSO (or DMSO only), and 0.02 mL of 100 nM MMP-1 or 4 nM MMP-2 or 300 nM MMP-3 were added, mixed, and the vial was held at 37°C for 5-180 minutes. Then 0.02 mL of 0.2 mM substrate was added, mixed and transferred into a pre-heated cell. The sample was allowed to equilibrate in the cuvette for 3-5 min at 37°C against small changes in temperature and changes in the enzyme-inhibitor equilibria related to addition of substrate. After that, the linear increase of fluorescence was monitored over 3-5 min, and the slope (V_0 or V_s) was obtained. Inhibitor concentrations were varied to collect data over V_s/V_0 ratio ranging from 0.05 to 0.95. The values of K_i^* were calculated by nonlinear weighted regression to the tight-binding equation according to Morrison and Walsh above:

$$V_s/V_0 = [1/(2 \times Et)] \times \text{SQR}[(K_i^* + It - Et)^2 + 4 \times K_i^* \times Et] - (K_i^* + It - Et)$$

wherein Et is the total enzyme concentration, and It is the total inhibitor concentration.

The following Table I lists K_i^* obtained for some representative examples:

TABLE 1

K_i at steady state (K_i^*) for inhibition of MMP-1, MMP-2 and MMP-3 (all micromolar)

Example compound	MMP-1	MMP-2	MMP-3
Example 2	1.1	2.8	13
Example 3	0.088	0.29	2.5
Example 5	0.061	0.31	0.94
Example 7	0.070	0.25	0.72

	<u>Example compound</u>	<u>MMP-1</u>	<u>MMP-2</u>	<u>MMP-3</u>
	Example 10	0.18	0.34	2.2
	Example 12	0.041	0.19	0.38
	Example 15	17	0.63	6.5
5	Example 16	6.4	0.094	2.4
	Example 17	9.4	0.61	4.3
	Example 18	0.85	0.079	0.86
	Example 19	4.5	0.23	1.1
	Example 20	2.0	0.04	0.39
10	Example 22	0.015	0.067	0.73
	Example 23	0.003	0.016	0.145
	Example 24	0.17	0.40	11
	Example 25	0.076	0.19	2.3
	Example 26	0.071	0.10	0.60
15	Example 27	0.0065	0.02	0.24
	Example 28	7.5	0.60	10
	Example 30	0.0036	0.025	0.27
	Example 32	2.6	0.037	0.50
	Example 33	>10	0.46	6.6
20	Example 34	2.4	0.021	1.4
	Example 35	0.0029	0.008	0.031
	Example 37	0.0018	0.0026	0.035
	Example 39	0.76	0.09	0.806
	Example 42	1.4	0.016	0.76
25	Example 44	0.0089	0.0023	0.21
	Example 47	0.0046	0.028	0.14
	Example 50	0.026	0.058	0.49
	Example 61	0.0078	0.066	not determined

Some of the compounds of formula (I) were also shown to possess high activity at inhibiting the release of TNF of several different cell lines, under different stimulation conditions. For example, the following cell-based assay was used to assess such activity:

CELLULAR ASSAY (Protocol B)

THP-1 cells, cultured in RPMI 1640 supplemented with 10% FCS, were distributed into 24-

well plates, 1 mL of a suspension of 1×10^6 cells/mL in each well. Compounds to be tested, dissolved in DMSO and diluted with the culture medium (1% final DMSO concentration) were added. Plates were incubated for 30 min at 37 °C in 5% CO₂, and lipopolisaccharide (LPS 0111:B4, 5 µg/mL) was added as a stimulant. After a further 4 h incubation, cells were harvested, centrifuged (2,000 rpm, 7 min), and the supernatant was collected and frozen (-20 °C) until analysis. Analysis was run by classical ELISA methodology (monoclonal anti-TNF- α antibody, rabbit capture polyclonal antibody, and peroxidated anti-rabbit antibody). Dichloroisocoumarin was used as a standard. As anticipated above, low aqueous solubility, metabolic instability (high clearance) and poor oral bioavailability is a common problem for the "peptide-based" hydroxamates of the prior art. In these respects, the compounds of the present invention are usually characterised by a superior profile. As an example, Table 2 reports the solubility in water for some representative compounds, detailed in the Examples.

TABLE 2

Aqueous solubility (mg/mL, 25 °C) of some representative compounds

Example compound	Conditions ¹	Solubility
Example 2	A	>4.5
Example 3	A	>9.5
Example 5	B	>6.0
Example 7	A	4.0
Example 13	A	6.71
Example 14	B	>14
Example 16	B	>9.6
Example 18	B	>9.6
Example 19	A	7.5
Example 22	B	>8.2
Example 27	B	7.1

(1) A: pH 7.4 phosphate buffer, from excess solid, after 5 min stirring; B: saline, from excess solid, after 5 min stirring

As a further example, Table 3 reports clearance and AUC values measured after intravenous administration to rats of a 10-15 mg bolus dose of some representative compounds of the present invention.

TABLE 3**PK parameters in rats (iv route, 10-15 mg/kg) of some representative compounds**

Example compound	Clearance (mL/min/kg)	AUC ($\mu\text{g}\cdot\text{min}/\text{mL}$) ¹
Example 10	24	666
5 Example 12	39	395
Example 19	27	563
Example 27	23	679

(1) Compounds were administered at doses ranging from 10 to 15 mg/kg. Data are referred to a
 10 15 mg/kg dose to allow comparison.

As an additional example, Table 4 reports the maximum plasma concentration (C_{max}) and oral
 bioavailability (%F; calculated from dose-normalised ratio of oral to i.v. mean AUC values) after
 oral administration to rats or cynomolgus monkeys of a 10-15 mg single dose of some
 15 representative compounds of the present invention.

TABLE 4**PK parameters (oral route, 10-15 mg/kg) of some representative compounds**

Example compound	Animal species	C_{max} (ng/mL) ¹	%F
Example 10	rat	2120	28
20 Example 12	rat	2440	21
Example 22	rat	5487	62
Example 27	rat	7908	58
Example 27	cynomolgus monkey	3132	34
Example 50	rat	3146	50
25 Example 50	cynomolgus monkey	49516	70

(1) Compounds were administered at doses ranging from 10 to 15 mg/kg. Data are referred to a
 15 mg/kg dose to allow comparison.

30 Compounds of formula (I) can be used in human or veterinary medicine in the form of
 pharmaceutical preparations which contain them in association with a compatible
 pharmaceutical carrier material. Thus, a distinct aspect of the present invention is the
 preparation of pharmaceutical compositions carrying a compound of formula (I) as active

ingredient, and a method of management (i.e. treatment or prophylaxis) of diseases or conditions mediated in humans and warm blood animals by MMPs and/or TACE, which method comprises administering an effective amount of a compound of formula (I) above, or a pharmaceutically acceptable salt thereof.

5 In particular, the compounds of formula (I) can be administered:

A) Orally, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known in the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, maize starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and adsorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example, peanut oil, liquid paraffin, or olive oil. Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example, sodium carboxymethylcellulose, methylcellulose, hydroxy propylmethylcellulose, sodium alginate, polyvinylpyrrolidone gum tragacanth and gum acacia; dispersing or wetting agents may be naturally-occurring phosphatides, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyoxyethylene

10
15
20
25
30

sorbitan monooleate. The said aqueous suspensions may also contain one or more preservatives, for example, ethyl or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, or one or more sweetening agents, such as sucrose or saccharin. Oily suspension may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, 5 olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents, such as those set forth above, and flavouring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an antioxidant such as ascorbic acid. Dispersible powders and granules suitable for preparation of an aqueous suspension 10 by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, a suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present. The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase 15 may be a vegetable oil, for example olive oil or arachis oils, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan mono-oleate, and condensation products of the said partial esters with ethylene oxide, for 20 example polyoxyethylene sorbitan monooleate. The emulsion may also contain sweetening and flavoring agents. Syrups and elixirs may be formulated with sweetening agents, for example glycerol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents;

B) Parenterally, either subcutaneously, or intravenously, or intramuscularly, or intrasternally, or by 25 infusion techniques, in the form of sterile injectable aqueous or oleaginous suspensions. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butane diol. Among the 30 acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition fatty acids such as oleic acid find use in the preparation of

injectables;

C) By inhalation, in the form of aerosols or solutions for nebulizers;

D) Rectally, in the form of suppositories prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperature but liquid at the rectal temperature and will
5 therefore melt in the rectum to release the drug. Such materials are cocoa butter and poly-ethylene glycols;

E) Topically, in the form of creams ointments, jellies, solutions or suspensions.

Daily doses are in the range of about 0.1 to about 50 mg per kg of body weight, according to the activity of the specific compound, the age, weight and conditions of the subject to be treated, the
10 type and severity of the disease, and the frequency and route of administration; preferably, daily dosage levels for humans are in the range of 10 mg to 2 g. The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. For example, a formulation intended for the oral administration to humans, may contain from 5 mg to 2 g of active agent
15 compounded with an appropriate and convenient amount of carrier material which may vary from about 5 to about 95 percent of the total composition. Dosage unit forms will generally contain between from about 5 mg to about 1 g of active ingredient.

Pharmaceutical compositions containing a compound of formula (I) can be used in medicine for the treatment of disease states characterised by a systemic or local imbalance of active
20 MMPs and their natural inhibitors. The rationale for the use of MMP inhibitors in medicine has been illustrated above, and is well described in the recent literature; see, for example, D.E. Levy & A.M. Ezrin, "Matrix Metalloproteinase Inhibitor Drugs", in: Emerging Drugs: The Prospect for Improved Medicines, Chapter Ten (pp 205-230), Ashley Publications Ltd., 1997. According to this rationale, and proofs of concept already established with other MMP
25 inhibitors, the compounds of the present invention can be used, in particular, for the treatment of:

- inflammatory and autoimmune diseases, especially rheumatoid arthritis, osteoarthritis, bone resorption, periodontal disease, multiple sclerosis, inflammatory bowel diseases;
- cancer, including both tumor growth and tumor invasion by secondary metastases, with
30 particular reference to breast cancer, small cell lung cancer, non-small cell lung cancer, glioblastoma, prostate cancer, ovarian cancer, gastric and esophageal cancers, pancreatic cancer, colorectal tumors, and bony metastases;
- angiogenic disorders, especially diabetic retinopathies and macular diseases;

- cardiovascular diseases, especially congestive heart failure and vascular restenosis;
- soft and osseous tissue diseases, including ocular inflammation, corneal or tissue ulceration, wound healing;
- other disorders in which either MMPs or abnormal release of TNF- α is implicated, in particular shock syndromes, transplant rejection, cachexia, anorexia.

The present invention also includes the use of compounds of formula (I), for the treatment of any of the diseases above, as adjuncts to other conventional treatments; for example, together with anti-inflammatory or immunosuppressive drugs for the treatment of rheumatoid arthritis and multiple sclerosis, and together with cytotoxic or cytostatic drugs for the treatment of tumoral diseases.

The following examples are meant to illustrate the present invention, without limiting it.

EXAMPLE 1

(3S-tert-Butoxycarbonylamino-4-hydroxy-2R-isobutyl)succinyl-piperidinamide

AZETIDINONE ROUTE

Step A

Piperidine (1.6 mL) was added to a solution of 4S-benzyloxycarbonyl-1-tert-butoxycarbonyl-3R-isobutylazetidin-2-one (3.8 g; see Preparation A) in acetonitrile (15 mL). After three hours stirring at room temperature, the solvent was removed in vacuo and the residue, dissolved in EtOAc, was sequentially washed with saturated aqueous NH_4Cl and brine. Drying over Na_2SO_4 , evaporation and flash chromatography over silica gel (n-hexane/EtOAc) afforded (4-benzyloxy-3S-tert-butoxycarbonylamino-2R-isobutyl)succinyl-piperidinamide as a yellow oil (3.6 g). FT-IR (CHCl_3) 3433 br (NH), 1753 (ester CO), 1714 (carbamate CO), 1626 (amide CO) cm^{-1} . ^1H -NMR (200 MHz, CDCl_3) 0.89 and 0.91 (two d, 6 H, $J = 6.6$ Hz); 1.42 (s, 9 H); 1.2-1.8 (m, 9 H); 3.1-3.5 (m, 5 H); 4.48 (dd, 1 H, $J = 4.1$ and 10.0 Hz); 5.06 and 5.19 (two d, 2 H, $J = 12.4$ Hz); 6.45 (d, 1 H, $J = 10.0$ Hz); 7.32 (m, 5 H) ppm.

Step B

A mixture of (4-benzyloxy-3S-tert-butoxycarbonylamino-2R-isobutyl)succinyl-piperidinamide (500 mg) and 10% Pd/C (100 mg) in EtOH (20 mL) was exposed to a hydrogen atmosphere for 3 hours. The catalyst was removed by filtration (Celite filter aid), washed with additional ethanol, and the solvent was removed in vacuo to leave the title compound (390 mg) as a colourless oil. ^1H -NMR (200 MHz, CDCl_3) 0.89 and 0.90 (two d, 6 H, $J = 6.1$ Hz) 1.44 (s, 9 H); 1.4-1.8 (m, 9 H); 3.60 (m, 5 H); 4.29 (dd, 1 H, $J = 1.9$ and 4.9 Hz);

5.92 (d, 1 H, J= 4.9 Hz) ppm. FAB-MS 379 (MNa)⁺, 357 (MH)⁺, 301, 257, 211, 112, 86, 84, 57 m/z.

ASPARTIC ROUTE

Starting from N-tert-butoxycarbonyl-3R-isobutyl-L-aspartic acid 1-benzyl ester-4-piperidin-
5 amide, obtained as described in Preparation H, which is (4-benzyloxy-3S-tert-butoxycarbonylamino-2R-isobutyl)succinyl-piperidin-amide, identical with the compound obtained as in step A above, and following the same procedure described in step B above, the title product was obtained, identical with the one described above.

EXAMPLE 2

10 **(3S-tert-Butoxycarbonylamino-4-hydroxyamino-2R-isobutyl)succinyl-piperidinamide**

Step A

(3S-tert-Butoxycarbonylamino-4-hydroxy-2R-isobutyl)succinyl-piperidinamide (390 mg),
obtained as indicated in Example 1, was dissolved in CH₃CN (20 mL) and treated with O-benzylhydroxylamine hydrochloride (230 mg) and N-methylmorpholine (0.33 ml). After 10
15 min, TBTU (O-1H-benzotriazol-1-yl-N,N,N',N'-tetramethyluronium tetrafluoroborate; 465 mg) was added to the solution, and the mixture was let stir overnight at room temperature. The solvent was removed in vacuo and the residue, dissolved in CH₂Cl₂, was washed with 4% aqueous NaHCO₃, 2% HCl and brine. The organic layer was dried over Na₂SO₄ and evaporated. The crude product was purified by flash chromatography on silica gel (n-hexane/EtOAc) to afford (4-benzyloxyamino-3S-tert-butoxycarbonylamino-2R-isobutyl)-
20 succinyl-piperidinamide as an oil (500 mg). H¹- NMR (200 MHz, CDCl₃) 0.90 and 0.92 (two d, 6 H, J= 6.4 Hz); 1.57 (s, 9 H); 1.3-1.7 (m, 9 H); 4.32 (dd, 1 H, J= 3.4 and 8.2 Hz); 4.85 (s, 2 H); 6.73 (d, 1 H, J= 8.5 Hz); 7.37 (m, 5 H); 9.13 (s, 1 H) ppm. ESI-MS 500 (MK)⁺, 484 (Mna)⁺, 462 (MH)⁺, 384, 362, 283, 239, 211, 112 m/z.

25 **Step B**

A mixture of (4-benzyloxyamino-3S-tert-butoxycarbonylamino-2R-isobutyl)succinyl-piperidinamide (500 mg) and 10% Pd/C (100 mg) in EtOH (50 mL) was exposed to a hydrogen atmosphere for 1 hour. The catalyst was removed by filtration (Celite filter aid) and the solvent was evaporated to leave the title compound (300 mg) as a white solid. H¹- NMR
30 (200 MHz, CDCl₃) 0.90 and 0.93 (two d, 6 H, J= 6.1 Hz); 1.45 (s, 9 H); 1.2-1.8 (m, 9 H); 3.3-3.7 (m, 5 H); 4.43 (dd, 1 H, J= 3.4 and 8.3 Hz); 6.72 (d, 1 H, J= 8.3 Hz); 6.93 (br signal, 1 H); 9.31 (br signal, 1 H) ppm. FAB-MS 372 (MH)⁺, 316, 283, 272, 211, 112, 86, 84, 57 m/z.

EXAMPLE 3**(3S-Amino-4-hydroxyamino-2R-isobutyl)succinyl-piperidinamide**

(3S-tert-Butoxycarbonylamino-4-hydroxyamino-2R-isobutyl)succinyl-piperidinamide (170 mg; prepared as described in Example 2) was dissolved in 95% aqueous trifluoroacetic acid (3 mL), and the solution was stirred 2 hours at room temperature. Toluene was added and evaporated in vacuo (several times), thereby obtaining the title compound, trifluoroacetate salt, as a white solid (115 mg). In DMSO solution, the compound is present as a mixture of two rotamers (M, major; m, minor). H^1 -NMR (400 MHz, DMSO- d_6) 0.80 and 0.81 (two d, 6 H, J = 6.4 Hz); 1.1-1.6 (m, 9 H); 3.21 (m, 1 H); 3.40 (m, 4 H); 3.64 (d, 1 H, J = 7.3 Hz); 8.02 (br signal, 3 H); 9.32 (s, 1 H, M); 9.80 (br signal, 1 H, m); 10.69 (s, 1 H, m); 11.12 (br signal, 1 H, M) ppm. ESI-MS 294 (MNa) $^+$, 272 (MH) $^+$, 239, 112 m/z.

EXAMPLE 4**(3S-N,N-Dimethylamino-4-hydroxy-2R-isobutyl)succinyl-piperidinamide****Step A**

(4-Benzyloxy-3S-tert-butoxycarbonylamino-2R-isobutyl)succinyl-piperidinamide (prepared as described Example 1, step A; 1 g) was dissolved in 95% aqueous trifluoroacetic acid (5 mL), and the solution was stirred 2 hr at room temperature. Toluene was added and evaporated in vacuo, repeating the process several times, thereby obtaining (3S-amino-4-benzyloxy-2R-isobutyl)succinyl-piperidinamide (trifluoroacetate salt) as a yellow oil (1g).

Step B

Sodium cyanoborohydride (230 mg) was added to a solution of (3S-amino-4-benzyloxy-2R-isobutyl)succinyl-piperidinamide trifluoroacetate salt (700 mg) in MeOH (10 mL). After 10 min, 37% aqueous formaldehyde was added dropwise, and the solution was stirred for 1 hour at room temperature. The solvent was removed in vacuo and the residue was dissolved in EtOAc; the organic layer was washed with water and dried over Na_2SO_4 . Evaporation of the solvent left crude (4-benzyloxy-3S-N,N-dimethylamino-2R-isobutyl)succinyl-piperidinamide (552 mg). H^1 -NMR (200 MHz, $CDCl_3$) 0.80 and 0.82 (two d, 6 H, J = 6.6 Hz); 0.92 and 1.70 (two m, 2 H); 1.40 (m, 1 H); 1.6 (m, 6 H); 2.25 (s, 6 H); 3.30 (m, 1 H); 3.50 (m, 4 H); 3.56 (d, 1 H, J = 10.8 Hz); 5.15 and 5.20 (two d, 2 H, J = 12.2 Hz); 7.36 (m, 5 H) ppm.

Step C

(4-Benzyloxy-3S-N,N-dimethylamino-2R-isobutyl)succinyl-piperidinamide (550 mg), as the crude product from step B above, was dissolved in EtOH (15 mL). The resulting solution

was treated with 10% Pd/C (100 mg) and exposed to a hydrogen atmosphere for 4 hours. The catalyst was removed by filtration (Celite filter aid), washing with additional EtOH. The solvent was removed in vacuo to leave the title compound (400 mg) as a yellow oil. FT-IR (KBr) 3385 br (NH), 1622 br (CO) cm^{-1} . ESI/CID- MS 307 (MNa)⁺, 285 (MH)⁺, 239 (M-COOH)⁺, 184, 102 m/z.

EXAMPLE 5

(3S-N,N-Dimethylamino-4-hydroxyamino-2R-isobutyl)succinyl-piperidinamide

Step A

(3S-N,N-Dimethylamino-4-hydroxy-2R-isobutyl)succinyl-piperidinamide (460 mg), prepared as described in Example 4 (step B), was dissolved in CH₃CN (10 mL) and treated with O-benzylhydroxylamine hydrochloride (310 mg) and N-methylmorpholine (0.44 mL). After 10 min, TBTU (663 mg) was added to the solution, and the mixture was let stir overnight at room temperature. The solvent was removed in vacuo and the residue, dissolved in EtOAc, was washed with aqueous NaHCO₃ and brine. The organic layer was dried over Na₂SO₄ and evaporated to afford the crude product. Purification by flash chromatography on silica gel (n-hexane/EtOAc) afforded a white solid (230 mg), consisting of (4-benzyloxyamino-3S-N,N-dimethylamino-2R-isobutyl)succinyl-piperidinamide. FT-IR (KBr) 3191 br (NH), 1683 br (CO) cm^{-1} . ¹H-NMR (400 MHz, DMSO-d₆) 0.75 and 0.80 (two d, 6 H, J= 6.4 Hz); 0.9-1.6 (m, 9 H); 2.09 (s, 6 H); 2.97 (d, 1 H, J= 10.7 Hz); 3.30 (m, 1 H); 3.3-3.6 (m, 4 H); 4.79 and 4.82 (two d, 2 H, J= 12.0 Hz); 7.36 (m, 5 H); 11.13 (s, 1 H) ppm. ESI-MS 412 (MNa)⁺, 390 (MH)⁺, 305, 239, 207 m/z.

Step B

The material from step A above (230 mg) and 10% Pd/C (50 mg) in EtOH (10 mL) was exposed to a hydrogen atmosphere for 1 hour. The catalyst was removed by filtration (Celite filter aid) and the solvent was evaporated to leave the title compound as a white solid (170 mg). ¹H-NMR (400 MHz, DMSO-d₆) 0.77 and 0.81 (two d, 6 H, J= 6.7 Hz); 0.9-1.6 (m, 9 H); 2.13 (s, 6 H); 3.00 (d, 1 H, J= 10.8 Hz); 3.30 (m, 1 H); 3.3-3.7 (m, 4 H); 8.83 (s, 1 H); 10.51 (s, 1 H) ppm. ESI-MS 322 (MNa)⁺, 300 (MH)⁺, 239 (M-HONHCO), 117 m/z.

EXAMPLE 6

(4-Hydroxy-3S-(4-methoxyphenylsulfonyl)amino-2R-isobutyl)succinyl-piperidinamide

Step A

(3S-Amino-4-benzyloxy-2R-isobutyl)succinyl-piperidinamide trifluoroacetate salt (prepared as described in Example 4, step B; 660 mg) was dissolved in CH₂Cl₂ (100 mL). To this

solution, (4-methoxybenzenesulfonyl)chloride (350 mg) and triethylamine (0.5 mL) were added. After 24 hours, the solution was washed with saturated aqueous NH_4Cl , saturated aqueous NaHCO_3 and brine. The organic layer was dried over Na_2SO_4 and evaporated to afford a crude product, which was purified by flash chromatography over silica gel (n-hexane/EtOAc) to give (4-benzyloxy-3S-(4-methoxyphenylsulfonyl)amino-2R-isobutyl)succinyl-piperidinamide (340 mg) as a white solid. ^1H -NMR (200 MHz, CDCl_3) 0.86 and 0.90 (two d, 6 H, $J=6.1$ Hz) 1.2-1.8 (m, 9 H); 3.0-3.5 (m, 5 H); 3.81 (s, 3 H); 4.20 (dd, 1 H, $J=3.9$ and 9.8 Hz); 4.90 (s, 2 H); 6.72 (d, 1 H, $J=9.8$ Hz); 6.82 (d, 2 H, $J=9.0$ Hz); 7.30 (m, 5 H); 7.75 (d, 2 H, $J=9.0$ Hz) ppm. ESI-MS 555 (MH)⁺, 539 (MNa)⁺, 454, 112, 91 m/z.

10 Step B

The material from step A above (340 mg) and 10% Pd/C (70 mg) were suspended in a mixture of EtOH and THF (1:1; 30 mL) and stirred under a hydrogen atmosphere for 2 hours. The catalyst was removed by filtration (Celite filter aid) and the solvent was evaporated, to leave the title compound as a white solid (245 mg). ^1H -NMR (400 MHz, CDCl_3) 0.91 and 0.94 (two d, 6 H, $J=6.4$ Hz) 1.2-1.8 (m, 9 H); 3.5-3.8 (m, 6 H); 3.84 (s, 3 H); 6.00 (br signal, 1 H); 6.95 (d, 2 H, $J=9.0$ Hz); 7.71 (d, 2 H, $J=9.0$ Hz) ppm. ESI-MS 465 (MK)⁺, 449 (MNa)⁺, 427 (MH)⁺, 364, 196 m/z.

EXAMPLE 7

20 (4-Hydroxyamino-3S-(4-methoxyphenylsulfonyl)amino-2R-isobutyl)succinyl-piperidinamide

Step A

(4-Hydroxy-3S-(4-methoxyphenylsulfonyl)amino-2R-isobutyl)succinyl-piperidinamide (245 mg; Example 6) was dissolved in CH_3CN (20 mL) and treated with O-benzylhydroxylamine hydrochloride (110 mg) and N-methylmorpholine (0.16 mL). After 10 min, TBTU (220 mg) was added, and the mixture was stirred overnight at room temperature. The solvent was removed in vacuo and the residue was dissolved in EtOAc and washed with saturated aqueous NaHCO_3 , 2% aqueous HCl and brine. The organic layer was dried over Na_2SO_4 and evaporated to afford a crude product which was purified by flash chromatography on silica gel (n-hexane/EtOAc) to obtain a white solid (300 mg), consisting of (4-benzyloxyamino-3S-(4-methoxyphenylsulfonyl)amino-2R-isobutyl)succinyl-piperidinamide. ^1H -NMR (200 MHz, CDCl_3) 0.71 and 0.72 (two d, 6 H, $J=6.3$ Hz); 0.8-1.7 (m, 9 H); 3.3-3.6 (m, 5 H); 3.81 (dd, 1 H, $J=6.6$ Hz); 4.80 and 4.83 (two d, 2 H, $J=9.0$ Hz); 6.94 (d, 2 H, $J=9.0$ Hz); 7.14 (d, 1

H, J= 6.4 Hz); 7.39 (m, 5 H); 7.76 (d, 2 H, J= 9.0 Hz); 9.65 (br signal, 1 H) ppm. ESI/CID-MS 570 (MK)⁺, 554 (MNa)⁺, 469 (MH)⁺, 403, 381, 171, 112, 91 m/z.

Step B

The material from step A above (300 mg) and 10% Pd/C (100 mg) were suspended in EtOH (20 mL) and stirred under a hydrogen atmosphere for 3 hr at room temperature. The catalyst was removed by filtration (Celite filter aid), washing with additional EtOH. The solvent was removed in vacuo to give the title compound (210 mg) as white solid. H¹- NMR (400 MHz, CDCl₃) 0.70 and 0.71 (two d, 6 H, J= 6.4 Hz); 0.9-1.7 (m, 9 H); 3.3-3.7 (m, 5 H); 3.86 (s, 3 H); 3.89 (dd, 1 H, J= 3.0 and 6.6 Hz); 6.98 (d, 2 H, J= 9.0 Hz); 7.23 (d, 1 H, J= 6.6 Hz); 7.82 (d, 2 H, J= 9.0 Hz); 9.79 (br signal, 1 H) ppm. ESI-MS 480 (MK)⁺, 464 (MNa)⁺, 442 (MH)⁺, 409, 381, 171 m/z.

EXAMPLE 8

(3S-tert-Butoxycarbonylamino-4-hydroxy-2R-isobutyl)succinyl-morpholinamide

Step A

In a manner analogous to that described in Example 1 (step A), (3S-tert-butoxycarbonylamino-2R-isobutyl-4-(4-nitrobenzyloxycarbonyl)succinyl-morpholinylamide (190 mg) was obtained as a yellow oil starting from 1-tert-butoxycarbonyl-3R-isobutyl-4S-p-nitrobenzyloxycarbonylazetidin-2-one (200 mg) and morpholine (0.08 mL). FT-IR (CHCl₃) 3432 br (NH), 1750-1713 br (ester and carbamate CO), 1630 (amide CO) cm⁻¹. H¹- NMR (400 MHz, CDCl₃) 0.92 and 0.98 (two d, 6 H); 1.2-1.7 (m, 3 H); 1.44 (s, 9 H); 3.4-3.7 (m, 9 H); 4.58 (dd, 1 H, J= 4.1 and 10.0 Hz); 5.21 and 5.26 (two d, 2 H, J= 13.7 Hz); 6.36 (d, 1 H, J= 10.0 Hz); 7.50 (d, 2 H, J= 9.5 Hz); 8.20 (d, 2 H, J= 9.5 Hz) ppm.

Step B

In a manner analogous to that described in Example 1 (step B), from the material of step A above (500 mg), with 10% Pd/C (50 mg) and under a hydrogen atmosphere, the title compound was obtained (290 mg) as a white solid. FT-IR (KBr) 3411 br (NH,OH), 1750-1714 br (acid and carbamate CO), 1630 (amide CO) cm⁻¹. H¹- NMR (200 MHz, DMSO-d₆) 0.82 and 0.83 (two d, 6 H, J= 6.0 Hz); 1.2-1.6 (m, 3 H); 1.36 (s, 9 H); 3.2-3.6 (m, 9 H); 4.11 (dd, 1 H, J= 9.4 Hz); 6.41 (d, 1 H, J= 9.4 Hz); 12.80 (br signal, 1 H) ppm. FD-MS 358 (MH)⁺, 313 (MH-COOH)⁺, 259 m/z.

EXAMPLE 9

(3S-tert-Butoxycarbonylamino-4-hydroxyamino-2R-isobutyl)succinyl-morpholinamide

Step A

In a manner analogous to that described in Example 2 (step A), reaction of (3S-tert-butoxycarbonylamino-4-hydroxy-2R-isobutyl)succinyl-morpholinamide (290 mg) with O-benzylhydroxylamine hydrochloride (154 mg), N-methylmorpholine (0.22 mL) and TBTU (0.97 mg) afforded (4-benzyloxyamino-3S-tert-butoxycarbonylamino-2R-isobutyl)succinyl-morpholinamide (300 mg) as a white solid. H^1 -NMR (400 MHz, $CDCl_3$) 0.91 and 1.11 (two d, 6 H, $J = 6.2$ Hz); 1.3-1.6 (m, 3 H); 1.4 (s, 9 H); 3.4-3.8 (m, 9 H); 4.32 (dd, 1 H, $J = 3.2$ and 8.2 Hz); 4.86 (s, 2 H); 6.64 (d, 1 H, $J = 8.2$ Hz); 7.37 (m, 5 H); 9.12 (s, 1 H) ppm.

Step B

In a manner analogous to that described in Example 2 (step B), from the material of step A above (300 mg) and 10% Pd/C (60 mg), under a hydrogen atmosphere, the title compound was obtained as white solid (240 mg). H^1 -NMR (400 MHz, $CDCl_3$) 0.90 and 0.92 (two d, 6 H, $J = 6.7$ Hz); 1.2-1.7 (m, 3 H); 1.45 (s, 9 H); 3.3-3.8 (m, 8 H); 3.55 (m, 1 H); 4.43 (m, 1 H); 6.63 (d, 1 H, $J = 8.2$ Hz); 7.20 (br signal, 1 H); 9.34 (br signal, 1 H) ppm. FAB-MS 374 (MH)⁺, 318, 285, 274, 213, 114, 88, 57 m/z.

EXAMPLE 10

(3S-Amino-4-hydroxyamino-2R-isobutyl)succinyl-morpholinamide

In a manner analogous to that described in Example 3, reaction of (3S-tert-butoxycarbonylamino-4-hydroxyamino-2R-isobutyl)succinyl-morpholinamide (100 mg) with 95% aqueous trifluoroacetic acid afforded the title compound, trifluoroacetate salt, as a white solid (100 mg). In DMSO solution, the compound is present as a mixture of two rotamers (M, major; m, minor). H^1 -NMR (400 MHz, DMSO- d_6) 0.78 and 0.79 (two d, 6 H, $J = 6.4$ Hz); 1.15-1.5 (two m, 2 H); 1.35 (m, 1 H); 3.1-3.7 (m, 10 H); 8.10 (br signal, 3 H); 9.35 (s, 1 H, M); 9.80 (s, 1 H, m); 10.72 (s, 1 H, m); 11.13 (br signal, 1 H, M) ppm. FAB-MS 274 (MH)⁺, 213 (M-HONHCO)⁺, 114, 89, 88 m/z.

EXAMPLE 11

(4-Hydroxy-2R-isobutyl-3S-(4-toluenesulfonyl)amino)succinyl-morpholinamide

Step A

3R-Isobutyl-4S-p-nitrobenzyloxycarbonyl-1-(4-toluenesulfonyl)-azetidin-2-one (300 mg; see Preparation B) was dissolved in dry DMF (15 mL). To this solution, morpholine (0.11 mL) and sodium azide (30 mg) were sequentially added. After overnight stirring at room temperature, the solvent was partially removed in vacuo and the residue, taken up in EtOAc, was sequentially washed with saturated aqueous NH_4Cl and brine. Drying over Na_2SO_4 , evaporation, and flash chromatography over silica gel (n-hexane/EtOAc) afforded (2R-

isobutyl-4-p-nitrobenzyloxycarbonyl-3S-(4-toluenesulfonyl)amino)succinyl-morpholinamide (290 mg) as a white solid. FT-IR (CHCl₃) 3410 br (NH), 1736 (ester CO), 1674 (amide CO) cm⁻¹. H¹- NMR (400 MHz, CDCl₃) 0.80 and 0.88 (two d, 6 H, J= 6.4 Hz); 1.2-1.6 (m, 3 H); 2.36 (s, 3 H); 3.3-3.7 (m, 9 H); 4.27 (dd, 1 H, J= 4.1 and 9.7 Hz); 4.98 and 5.08 (two d, 2 H, J= 13.5 Hz); 6.67 (d, 1 H, J= 9.7 Hz); 7.19 (d, 2 H, J= 8.2 Hz); 7.36 (d, 2 H, J= 8.8 Hz); 7.71 (d, 2 H, J= 8.2 Hz); 8.18 (d, 2 H, J= 8.8 Hz) ppm.

Step B

In a manner analogous to that described in Example 1 (step B), from the material of step A above (290 mg), with 10% Pd/C (50 mg) under a hydrogen atmosphere, the title compound was obtained as an amorphous solid (200 mg). H¹- NMR (400 MHz, CDCl₃) 0.90 and 0.93 (two d, 6 H); 1.5 (m, 3 H); 2.40 (s, 3 H); 3.5-3.9 (m, 10 H); 6.36 (d, 1 H, J= 3 Hz); 7.29 (d, 2 H, J= 8.1 Hz); 7.66 (d, 2 H, J= 8.1 Hz) ppm. FAB-MS 413 (MH)⁺, 367 (MH-COOH)⁺, 259, 213, 155, 114, 88 m/z.

EXAMPLE 12

(4-Hydroxyamino-2R-isobutyl-3S-(4-toluenesulfonyl)amino)succinyl-morpholinamide

Step A

In a manner analogous to that described in Example 7 (step A), reaction of (4-hydroxy-2R-isobutyl-3S-(4-toluenesulfonyl)amino)succinyl-morpholinamide (200 mg) with O-benzyl-hydroxylamine hydrochloride (93 mg), N-methylmorpholine (0.13 mL) and TBTU (186 mg) afforded (4-benzyloxyamino-2R-isobutyl-3S-(4-toluenesulfonyl)amino)succinyl-morpholinamide (176 mg) as a white solid. H¹- NMR (400 MHz, CDCl₃) 0.67 and 0.68 (two d, 6 H, J= 6.4 Hz); 0.8-1.2 (m, 3 H); 2.42 (s, 3 H); 3.3-3.8 (m, 10 H); 4.80 and 4.86 (two d, 2 H, J= 10.8 Hz); 7.04 (d, 1 H, J= 6.4 Hz); 7.30 (d, 2 H, J= 8.6 Hz); 7.39 (m, 5 H); 7.72 (d, 2 H, J= 8.6 Hz); 9.53 (s, 1 H) ppm.

Step B

In a manner analogous to that described in Example 7 (step B), from the material of step A above (170 mg) with 10% Pd/C (50 mg) under hydrogen atmosphere, the title compound was obtained as a white solid. H¹- NMR (400 MHz, DMSO-d₆) 0.67 and 0.70 (two d, 6 H, J= 6.8 Hz); 1.1-1.5 (m, 3 H); 2.30 (s, 3 H); 3.0-3.6 (m, 9 H); 3.72 (m, 1 H); 7.25 (d, 2 H, J= 8.1 Hz); 7.53 (d, 2 H, J= 8.1 Hz); 7.89 (d, 1 H, J= 9.4 Hz); 8.82 (s, 1 H); 10.74 (s, 1 H) ppm. FAB-MS 428 (MH)⁺, 371, 367 (M-HONHCO)⁺, 241, 129, 114, 57 m/z.

EXAMPLE 13

(3S-tert-Butoxycarbonylamino-4-hydroxy-2R-(4-methoxy)phenpropyl)succinyl-morpholinamide

Step A

In a manner analogous to that described in Example 1 (step A), reaction of 4S-benzyloxycarbonyl-1-tert-butoxycarbonyl-3R-(4-methoxy)phenpropyl)azetidin-2-one (600 mg; see Preparation C) with morpholine (0.23 mL) afforded (4-benzyloxy-3S-tert-butoxycarbonylamino-2R-(4-methoxy)phenpropyl)succinyl-morpholinamide as a yellow oil (700 mg). H^1 -NMR (400 MHz, DMSO- d_6) 1.41 (s, 9 H); 1.4-1.8 (m, 4 H); 2.54 (m, 2 H); 3.1-3.5 (m, 8 H); 3.25 (m, 1 H); 3.76 (s, 3 H); 4.53 (dd, 1 H, J = 3.8 and 9.8 Hz); 5.02 and 5.19 (two d, 2 H, J = 12.4 Hz); 6.28 (d, 1 H, J = 9.8 Hz); 6.78 (d, 2 H, J = 8.6 Hz); 7.03 (d, 2 H, J = 8.6 Hz); 7.31 (s, 5 H) ppm.

Step B

In a manner analogous to that described in Example 1 (step B), from the material of step A above (700 mg) with 10% Pd/C (100 mg), under a hydrogen atmosphere, the title compound was obtained as a white solid (600 mg). FT-IR (KBr) 3400 br (OH, NH), 1710 (acid CO), 1700 (carbamate CO), 1650 (amide CO) cm^{-1} . H^1 -NMR (400 MHz, $CDCl_3$) 1.43 (s, 9 H); 1.6 (m, 4 H); 2.52 (m, 2 H); 3.3-3.8 (m, 9 H); 3.76 (s, 3 H); 4.32 (dd, 1 H, J = 1.7 and 5.0 Hz); 6.04 (d, 1 H, J = 5.0 Hz); 6.80 (d, 2 H, J = 8.5 Hz); 7.02 (d, 2 H, J = 8.5 Hz) ppm. ESI-MS 473 (MNa) $^+$, 451 (MH) $^+$, 395, 351 (M-BOC) $^+$ m/z.

EXAMPLE 14

(3S-Amino-4-hydroxy-2R-(4-methoxy)phenpropyl)succinyl-morpholinamide

(3S-tert-Butoxycarbonylamino-4-hydroxy-2R-(4-methoxy)phenpropyl)succinyl-morpholinamide (obtained as described in Example 13; 50 mg) was dissolved in 95% aqueous trifluoroacetic acid (2 mL), and the solution was stirred 2 hours at room temperature. Toluene was added and evaporated in vacuo to afford the title compound as a white solid (58 mg). H^1 -NMR (400 MHz, DMSO- d_6) 1.4-1.6 (m, 4 H); 2.48 (m, 2 H); 3.2-3.6 (m, 9 H); 3.70 (s, 3 H); 3.91 (d, 1 H, J = 4.1 Hz); 6.82 (d, 2 H, J = 8.5 Hz); 7.09 (d, 2 H, J = 8.5 Hz); 8.1 (br signal, 2 H) ppm. ESI-MS 373 (MNa) $^+$, 351 (MH) $^+$, 305 (M-COOH) $^+$, 236, 191 m/z.

EXAMPLE 15

(3S-tert-Butoxycarbonylamino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-morpholinamide

Step A

In a manner analogous to that described in Example 2 (step A), reaction of (3S-tert-butoxycarbonylamino-4-hydroxy-2R-(4-methoxy)phenpropyl)succinyl-morpholinamide (310 mg; obtained as described in Example 13) with O-benzylhydroxylamine hydrochloride (132 mg), N-methylmorpholine (0.2 mL) and TBTU (263 mg) afforded (4-benzyloxyamino-3S-
5 tert-butoxycarbonylamino-2R-(4-methoxy)phenpropyl)succinyl-morpholinamide as a white solid (300 mg). FT-IR (KBr) 3208 br (NH), 1710-1667 br (CO), 1632 (amide CO) cm^{-1} . ^1H -NMR (400 MHz, CDCl_3) 1.38 (s, 9 H); 1.5-1.8 (m, 4 H); 2.52 (m, 2 H); 3.3-3.8 (m, 9 H); 3.76 (s, 3 H); 4.33 (dd, 1 H, $J=3.0$ and 8.0 Hz); 4.83 (s, 2 H); 6.63 (d, 1 H, $J=8.0$ Hz); 6.79 (d, 2 H, $J=8.5$ Hz); 7.03 (d, 2 H, $J=8.5$ Hz); 7.35 (m, 5 H); 9.14 (s, 1 H) ppm. ESI-MS 578
10 $(\text{MNa})^+$, 555 $(\text{MH})^+$, 500, 456, 377, 305 m/z.

Step B

In a manner analogous to that described in Example 2 (step B), from the material of step A above (300 mg), with 10% Pd/C (50 mg) under a hydrogen atmosphere, the title compound was obtained as pinkish solid (250 mg). FT-IR (CHCl_3) 3254 br (NH,OH), 1712-1614 br
15 (CO) cm^{-1} . ^1H -NMR (400 MHz, CDCl_3) 1.44 (s, 9 H); 1.4-1.8 (m, 4 H); 2.54 (m, 2 H); 3.3-3.8 (m, 9 H); 3.78 (s, 3 H); 4.46 (dd, 1 H, $J=3.0$ and 8.1 Hz); 6.63 (d, 1 H, $J=8.1$ Hz); 6.81 (d, 2 H, $J=8.5$ Hz); 7.05 (d, 2 H, $J=8.5$ Hz); 7.6 (br signal, 1 H); 8.43 (br signal, 1 H) ppm. ESI-MS 488 $(\text{MNa})^+$, 466 $(\text{MH})^+$, 410, 377, 366, 305 m/z.

EXAMPLE 16

20 (3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-morpholinamide

In a manner analogous to that described in Example 3, reaction of (3S-tert-butoxycarbonylamino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-morpholinylamide (200 mg; obtained as described in Example 15) with 95% aqueous trifluoroacetic acid (2 mL) afforded the title compound, trifluoroacetate salt, as a white solid (154 mg). The compound
25 is present in DMSO solution as a mixture of two rotamers (M, major; m, minor). ^1H -NMR (400 MHz, $\text{DMSO}-d_6$) 1.42 (m, 4 H); 2.41 (m, 2 H); 3.14 (m, 1 H); 3.2-3.6 (m, 8 H); 3.66 (s, 3 H); 3.72 (m, 1 H); 6.79 (d, 2 H, $J=8.5$ Hz); 7.03 (d, 2 H, $J=8.5$ Hz); 8.0 (br signal, 3 H, m); 8.15 (br signal, 3 H, M); 9.39 (br signal, 1 H, M); 9.79 (br signal, 1 H, m); 10.72 (s, 1 H, m); 11.15 (s, 1 H, M) ppm. ES/CID-MS 366 $(\text{MH})^+$, 305 (M-HONHCO), 218, 121, 114,
30 88 m/z.

EXAMPLE 17

(3S-tert-Butoxycarbonylamino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-piperidinamide

Step A

In a manner analogous to that described in Example 1 (step A), reaction of 4S-benzoyloxycarbonyl-1-tert-butoxycarbonyl-3-(4-methoxy)phenpropyl)azetidin-2-one (350 mg; see Preparation D) with piperidine (0.15 mL) afforded (4-benzoyloxycarbonyl-3S-tert-butoxycarbonylamino-2R-(4-methoxy)phenpropyl)succinyl-piperidinamide as a yellow oil (400 mg).
5 FT-IR (CHCl₃) 3400 br (NH), 1730 (ester CO), and 1641 br (carbamate and amide CO) cm⁻¹. ¹H¹- NMR (200 MHz, CDCl₃) 1.2-1.8 (m, 10 H); 1.55 (s, 9 H); 2.55 (m, 2 H); 3.1-3.4 (m, 5 H); 3.78 (s, 3 H); 4.52 (dd, 1 H, J= 4.1 Hz); 5.06 and 5.18 (two d, 2 H, J= 12.4 Hz); 6.43 (d, 1 H, J= 9.8 Hz); 6.81 (d, 2 H, J= 8.5 Hz); 7.06 (d, 2 H, J= 8.5 Hz); 7.32 (s, 5 H) ppm.
10 ESI-MS 561 (MNa)⁺, 539 (MH)⁺, 483, 439 m/z.

Step B

In a manner analogous to that described in Example 1 (step B), from the material of step A above (250 mg), with 10% Pd/C (50 mg) under a hydrogen atmosphere, (3S-tert-butoxycarbonylamino-4-hydroxy-2R-(4-methoxy)phenpropyl)succinyl-piperidinamide was
15 obtained as a colourless oil (200 mg).

Step C

In a manner analogous to that described in Example 2 (step A), reaction of (3S-tert-butoxycarbonylamino-4-hydroxy-2R-(4-methoxy)phenpropyl)succinyl-piperidinamide (200 mg) with O-benzylhydroxylamine hydrochloride (85 mg), N-methylmorpholine (0.12 mL) and TBTU
20 (172 mg) afforded (4-benzoyloxyamino-3S-tert-butoxycarbonylamino-2R-(4-methoxy)phenpropyl)succinyl-piperidinamide (240 mg) as a white solid. FT-IR (CHCl₃) 3236 br (NH), 1705 (benzylhydroxamate CO), 1678 (carbamate CO), 1614 (amide CO) cm⁻¹. ¹H¹- NMR (400 MHz, CDCl₃) 1.39 (s, 9 H); 1.4-1.8 (m, 10 H); 2.54 (m, 2 H); 3.3-3.6 (m, 5 H); 3.78 (s, 3 H); 4.35 (dd, 1 H, J= 3.5 and 8.5 Hz); 4.85 (s, 2 H); 6.73 (d, 1 H, J= 8.5 Hz); 6.81
25 (d, 2 H, J= 8.5 Hz); 7.05 (d, 2 H, J= 8.5 Hz); 7.38 (m, 5 H); 9.18 (s, 1 H) ppm. ESI-MS 576 (MNa)⁺, 554 (MH)⁺, 498, 454, 375, 303 m/z.

Step D

In a manner analogous to that described in Example 2 (step B), from the material from step C above (240 mg), with 10% Pd/C (50 mg), under a hydrogen atmosphere, the title compound
30 was obtained as a pinkish solid (175 mg). FT-IR (CHCl₃) 3255 br (OH,NH), 1710 (hydroxamic CO), 1665 (carbamate CO), 1612 (amide CO) cm⁻¹. ¹H¹- NMR (400 MHz, DMSO-d₆) 1.35 (s, 9 H); 1.3-1.6 (m, 10 H); 2.42 (m, 2 H); 3.2-3.4 (m, 5 H); 3.69 (s, 3 H); 4.02 (dd, 1 H, J= 6.2 and 9.1 Hz); 6.58 (d, 1 H, J= 9.1 Hz); 6.80 (d, 2 H, J= 8.5 Hz); 7.04

(d, 2 H, $J = 8.5$ Hz); 8.78 (br signal, 1 H); 10.67 (br signal, 1 H) ppm. ESI-MS 486 (MNa)⁺, 464 (MH)⁺, 408, 375, 364, 303 m/z.

EXAMPLE 18

(3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-piperidinamide

- 5 In a manner analogous to that described in Example 3, reaction of (3S-tert-butoxycarbonyl-amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-piperidinamide (125 mg; from Example 17) with 95% aqueous trifluoroacetic acid (2 mL) and work-up afforded the title compound, trifluoroacetate salt, as a white solid (90 mg). FT-IR (KBr) 3250 br (OH,NH), 1668 (hydroxamic CO), 1603 (amide CO) cm⁻¹. In DMSO solution, the compound is present
- 10 as a mixture of two rotamers (M, major; m, minor). H¹- NMR (400 MHz, DMSO-d₆) 1.2-1.6 (m, 10 H); 2.44 (m, 2 H); 3.17 (m, 1 H); 3.2-3.8 (m, 5 H); 3.70 (s, 3 H); 6.82 (d, 2 H, $J = 8.5$ Hz); 7.06 (d, 2 H, $J = 8.5$ Hz); 7.9 (br signal, 3 H); 9.32 (br signal, 1 H, M); 9.7 (br signal, 1 H, m); 10.7 (br signal, 1 H, m); 11.1 (br signal, 1 H, M) ppm. ESI/CID-MS 386 (MNa)⁺, 364 (MH)⁺, 303 (M-HONHCO)⁺, 112 m/z.

15 EXAMPLE 19

(3S-N,N-Dimethylamino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-morpholinamide

Step A

- In a manner analogous to that described in Example 4 (step A), reaction of the compound
- 20 described in Example 13, step A (600 mg), with 95% aqueous trifluoroacetic acid (5 mL) and work-up afforded (3S-amino-4-benzyloxy-2R-(4-methoxy)phenpropyl)succinyl-morpholinamide (500 mg), trifluoroacetate salt, as a yellow oil.

Step B

- In a manner analogous to that described in Example 4 (step B), reaction of the material from
- 25 step A above (350 mg) with 37% aqueous HCHO (10 mL) and NaCNBH₃ (100 mg) afforded (4-benzyloxy-3S-N,N-dimethylamino-2R-(4-methoxy)phenpropyl)succinyl-morpholinamide as a colourless oil (250 mg).

Step C

- In a manner analogous to that described in Example 4 (step C), reaction of the material from
- 30 step B above (250 mg) with 10% Pd/C (50 mg) under a hydrogen atmosphere afforded (3S-N,N-dimethylamino-4-hydroxy-2R-(4-methoxy)phenpropyl)succinyl-morpholinamide as a colourless oil (190 mg).

Step D

In a manner analogous to that described in Example 5 (step A), reaction of the material from step C above (190 mg) with O-benzylhydroxylamine hydrochloride (95 mg), N-methylmorpholine (0.15 mL) and TBTU (193 mg) afforded (4-benzyloxyamino-3S-N,N-dimethylamino-2R-(4-methoxy)phenpropyl)succinyl-morpholinamide (160 mg) as a colourless oil.

Step E

In a manner analogous to that described in Example 5 (step B), reaction of the material from step D above (160 mg) with 10% Pd/C (30 mg) under a hydrogen atmosphere afforded the title compound (130 mg) as a white solid. FT-IR (CHCl₃) 3229 br (OH, NH); 1659 (hydroxamic CO); 1614 (amide CO) cm⁻¹. H¹- NMR (200 MHz, DMSO-d₆) 1.0-1.5 (m, 4 H); 2.13 (s, 4 H); 2.40 (m, 2 H); 3.05 (d, 1 H, J= 10.7 Hz); 3.2-3.7 (m, 9 H); 3.69 (s, 3 H); 6.80 (d, 2 H, J= 8.8 Hz); 7.04 (d, 2 H, J= 8.8 Hz); 8.85 (br signal, 1 H); 10.50 (br signal, 1 H) ppm. ESI-MS 416 (MNa)⁺, 394 (MH)⁺, 333 (M-HONHCO)⁺ m/z.

EXAMPLE 20

(4-Hydroxyamino-3S-(4-methoxyphenylsulfonyl)amino-2R-(4-methoxy)phenpropyl)succinyl-morpholinamide

Step A

In a manner analogous to that described in Example 6 (step A), reaction of the material from Example 19, step A (270 mg), with (4-methoxybenzenesulfonyl)chloride (185 mg) and triethylamine (0.17 mL) afforded (4-benzyloxy-3S-(4-methoxyphenylsulfonyl)amino-2R-(4-methoxy)phenpropyl)succinyl-morpholinamide as a white solid (100 mg). H¹- NMR (200 MHz, CDCl₃) 1.5-1.9 (m, 4 H); 2.54 (m, 2 H); 3.12 (m, 1 H); 3.0-3.5 (m, 8 H); 3.78 and 3.82 (two s, 6 H); 4.25 (dd, 1 H, J= 3.9 and 9.8 Hz); 4.89 (s, 2 H); 6.52 (d, 1 H, J= 9.8 Hz); 6.80 (m, 4 H); 7.06 (d, 2 H, J= 8.8 Hz); 7.1-7.3 (m, 5 H); 7.76 (d, 2 H, J= 8.8 Hz) ppm. ESI/CID-MS 633 (MNa)⁺, 611 (MH)⁺, 546, 171, 88 m/z.

Step B

In a manner analogous to that described in Example 6 (step B), reaction of the material from step A above (100 mg) with 10% Pd/C (20 mg) under a hydrogen atmosphere afforded 90 mg of (4-hydroxy-3S-(4-methoxyphenylsulfonyl)amino-2R-(4-methoxy)phenpropyl)succinyl-morpholinamide was obtained as a colourless oil.

Step C

In a manner analogous to that described in Example 7 (step A), reaction of the material from step B above (90 mg) with O-benzylhydroxylamine (33 mg), N-methylmorpholine (0.05 mL)

and TBTU (65 mg) afforded (4-benzyloxyamino-3S-(4-methoxyphenylsulfonyl)amino-2R-(4-methoxy)phenpropyl)succinyl-morpholinamide (88 mg) as a white solid. H^1 -NMR (200 MHz, $CDCl_3$) 1.0-1.5 (m, 4 H); 2.32 (m, 2 H); 3.2-4.0 (m, 10 H); 3.78 and 3.84 (two s, 6 H); 4.79 and 4.83 (two d, 2 H, J = 11.0 Hz); 6.81 (d, 2 H, J = 8.7 Hz); 6.92 (m, 4 H); 7.39 (m, 5 H); 7.75 (d, 2 H, J = 8.8 Hz); 9.64 (s, 1 H) ppm. ESI-MS 648 (MNa)⁺, 626 (MH)⁺, 503, 475, 361 m/z.

Step D

In a manner analogous to that described in Example 7 (step B), reaction of the material from step C above (88 mg) with 10% Pd/C (20 mg) under a hydrogen atmosphere afforded the title compound as a white solid (70 mg). FT-IR ($CHCl_3$) 3266 br (OH,NH), 1678 br (hydroxamic CO), 1612 (amide CO) cm^{-1} . H^1 -NMR (400 MHz, DMSO- d_6) 1.0-1.5 (m, 4 H); 2.33 (m, 2 H); 2.97 (ddd, 1 H, J = 3.8, 9.8 and 10.7 Hz); 3.1-3.6 (m, 8 H); 3.68 and 3.78 (two s, 6 H); 3.78 (d, 1 H, J = 10.7 Hz); 6.79 (d, 2 H, J = 8.5 Hz); 7.00 (m, 4 H); 7.60 (d, 2 H, J = 9.0 Hz); 7.80 (br signal, 1 H); 8.84 (s, 1 H); 10.76 (br signal, 1 H) ppm. FAB-MS 536 (MH)⁺, 475 (M-HONHCO)⁺, 364, 305, 171, 121, 114, 88 m/z.

EXAMPLE 21

(3S-tert-Butoxycarbonylamino-2R-cyclopentylmethyl-4-hydroxyamino)succinyl-piperidinamide

Step A

In a manner analogous to that described in Example 1 (step A), reaction of 4S-benzyloxycarbonyl-1-tert-butoxycarbonyl-3R-cyclopentylmethylazetidin-2-one (1.25 g, obtained as described in Preparation D) with piperidine (0.65 mL) afforded (4-benzyloxy-3S-tert-butoxycarbonylamino-2R-cyclopentylmethyl)succinyl-piperidinamide (1.3 g) as a yellow oil. H^1 -NMR (400 MHz, DMSO- d_6) 0.9-1.7 (m, 17 H); 1.32 (s, 9 H); 3.2-3.4 (m, 5 H); 4.30 (dd, 1 H, J = 5.3 and 9.4 Hz); 5.07 and 5.13 (two d, 2 H, J = 12.6 Hz); 6.68 (d, 1 H, J = 9.4 Hz); 7.36 (m, 5 H) ppm. ESI-MS 495 (MNa)⁺, 473 (MH)⁺, 417, 373 m/z.

Step B

In a manner analogous to that described in Example 1 (step B), reaction of the material from step A above (500 mg) with 10% Pd/C (100 mg) under a hydrogen atmosphere afforded (3S-tert-butoxycarbonylamino-2R-cyclopentylmethyl-4-hydroxy)succinyl-piperidinamide as a colourless oil (410 mg).

Step C

In a manner analogous to that described in Example 2 (step A), reaction of (3S-tert-butoxycarbonylamino-2R-cyclopentylmethyl-4-hydroxy)succinyl-piperidinamide (410 mg) with O-benzylhydroxylamine hydrochloride (205 mg), N-methylmorpholine (0.29 mL) and TBTU (410 mg) afforded (4-benzyloxyamino-3S-tert-butoxycarbonylamino-2R-cyclopentyl-
5 methyl)succinyl-piperidinamide as a white solid (500 mg). ^1H -NMR (400 MHz, CDCl_3) 1.0 and 1.8 (m, 17 H); 1.37 (s, 9 H); 3.4-3.6 (m, 5 H); 4.32 (dd, 1 H, $J = 3.4$ and 8.5 Hz); 4.83 (s, 2 H); 6.73 (d, 1 H, $J = 8.5$ Hz); 7.37 (m, 5 H); 9.16 (s, 1 H) ppm. ESI-MS 510 (MNa^+), 488, 388, 309, 237 m/z.

Step D

10 In a manner analogous to that described in Example 2 (step B), reaction of the material from step C above (500 mg) with 10% Pd/C (100 mg) under a hydrogen atmosphere afforded the title compound was obtained as a pink solid (400 mg). ^1H -NMR (200 MHz, CDCl_3) 1.0 and 1.9 (m, 17 H); 1.45 (s, 9 H); 3.3-3.7 (m, 5 H); 4.45 (dd, 1 H, $J = 3.2$ and 8.3 Hz); 6.77 (d, 1 H, $J = 8.3$ Hz); 6.80 (br signal, 1 H); 9.32 (br signal, 1 H) ppm. ESI-MS 420 (MNa^+), 398
15 (MH^+), 342, 309, 298, 237 m/z.

EXAMPLE 22

(3S-Amino-2R-cyclopentylmethyl-4-hydroxyamino)succinyl-piperidinamide

Step A

A solution of 4S-benzyloxycarbonyl-1-tert-butoxycarbonyl-3R-cyclopentylmethylazetidin-2-one (4.6 g; obtained as described in Preparation D) in acetonitrile (100 mL) was treated with
20 piperidine (2 mL) and stirred at r.t. for 6 h. The mixture was concentrated *under vacuum* and the residue was poured into EtOAc/water. The organic layer was washed sequentially with 4% aqueous hydrochloric acid, brine, 4% aqueous hydrogen bicarbonate and brine, then dried over sodium sulfate and rotoevaporated. (4-Benzyloxy-3S-tert-butoxycarbonylamino-
25 2R-cyclopentylmethyl)succinyl-piperidinamide (5.6 g) was obtained as a yellow oil. ^1H -NMR (400 MHz, $\text{DMSO}-d_6$) 0.9-1.7 (m, 17 H); 1.32 (s, 9 H); 3.2-3.4 (m, 5 H); 4.30 (dd, 1 H, $J = 5.3$ and 9.4 Hz); 5.07 and 5.13 (two d, 2 H, $J = 12.6$ Hz); 6.68 (d, 1 H, $J = 9.4$ Hz); 7.36 (m, 5 H) ppm. ESI-MS 495 (MNa^+), 473 (MH^+), 417, 373 m/z.

Step B

30 A mixture of the compound from Step A above (5.6 g) and 5% Pd/C (1.2 g) in EtOH (200 mL) was exposed to a hydrogen atmosphere for 2 hours under stirring. After purging the mixture with nitrogen, the catalyst was removed by filtration over Celite, and the filtrate was

concentrated to dryness under reduced pressure, giving (3S-*tert*-butoxycarbonylamino-2R-cyclopentylmethyl-4-hydroxy)succinyl-piperidinamide (4.5 g) as a colourless oil.

Step C

The compound from Step B above (4.5 g) in acetonitrile (150 mL) was treated sequentially with O-benzylhydroxylamine hydrochloride (2.07 g), N-methylmorpholine (2.86 mL) and TBTU (4.17 g). The reaction mixture was stirred at room temperature for 3 hr, then concentrated *under vacuum* to about one third of its initial volume and partitioned between EtOAc and water. The upper layer was separated and washed sequentially with 2% aqueous hydrochloric acid, water, 4% aqueous hydrogen bicarbonate and brine. After drying (sodium sulfate), removal of the solvent afforded (4-benzyloxyamino-3S-*tert*-butoxycarbonylamino-2R-cyclopentylmethyl)succinyl-piperidinamide as a colourless oil (5.65 g). H^1 - NMR (400 MHz, $CDCl_3$) 1.0 and 1.8 (m, 17 H); 1.37 (s, 9 H); 3.4-3.6 (m, 5 H); 4.32 (dd, 1 H, $J = 3.4$ and 8.5 Hz); 4.83 (s, 2 H); 6.73 (d, 1 H, $J = 8.5$ Hz); 7.37 (m, 5 H); 9.16 (s, 1 H) ppm. ESI-MS 510 (MNa^+), 488, 388, 309, 237 m/z.

Step D

A mixture of the compound from Step C above (5.65 g) and 5% Pd/C (0.56 g) in ethanol (300 mL) was exposed to a hydrogen atmosphere for 1 h under stirring. The reaction mixture was purged with nitrogen, then filtered over Celite and eventually rotoevaporated to give (3S-*tert*-butoxycarbonylamino-2R-cyclopentylmethyl-4-hydroxyamino)succinyl-piperidinamide as a whitish solid (4.6 g). H^1 - NMR (200 MHz, $CDCl_3$) 1.0 and 1.9 (m, 17 H); 1.45 (s, 9 H); 3.3-3.7 (m, 5 H); 4.45 (dd, 1 H, $J = 3.2$ and 8.3 Hz); 6.77 (d, 1 H, $J = 8.3$ Hz); 6.80 (br signal, 1 H); 9.32 (br signal, 1 H) ppm. ESI-MS 420 (MNa^+), 398 (MH^+), 342, 309, 298, 237 m/z.

Step E

The compound from Step D above (4.6 g) was treated with 95% aqueous trifluoroacetic acid (40 mL) and let stand at r.t. for 45 minutes, after which time toluene (50 mL) was added, and the resulting mixture was concentrated to dryness under reduced pressure. The residue was purified by reverse-phase chromatography (LiChroprep RP C-18) eluting with acetonitrile-water mixtures (from 0:100 to 50:50) containing 1% TFA. Product containing fractions were collected and concentrated *in vacuo*. (3S-Amino-2R-cyclopentylmethyl-4-hydroxyamino)-succinyl-piperidinamide was thus obtained as a white powder (trifluoroacetate salt, 3.7 g; ca. 75% overall yield). In DMSO solution, the compound is present as a mixture of two rotamers (M, major; m, minor). H^1 - NMR (400 MHz, DMSO-

d_s) 0.9-1.8 (m, 17 H); 3.16 (m, 1 H); 3.40 (m, 4 H); 3.67 (m, 1 H); 7.95 (br signal, 3 H, m); 8.14 (br signal, 3 H, M); 9.33 (s, 1 H, M); 9.77 (s, 1 H, m); 10.69 (s, 1 H, m); 11.12 (s, 1 H, M) ppm. FAB-MS 298 (MH)⁺, 237 (M-HONHCO)⁺, 208, 112, 86, 84 m/z.

EXAMPLE 23

5 **(2R-Cyclopentylmethyl-4-hydroxyamino-3S-(4-methoxyphenylsulfonyl)amino)succinyl-piperidinamide**

Step A

In a manner analogous to that described in Example 6 (step A), reaction of (3S-amino-4-benzyloxy-2R-cyclopentylmethyl)succinyl-piperidinamide trifluoroacetate salt (630 mg) with
10 (4-methoxyphenylsulfonyl)chloride (400 mg) and triethylamine (0.78 mL) afforded (4-benzyloxy-2R-cyclopentylmethyl-3S-(4-methoxyphenylsulfonyl)amino)succinyl-piperidinamide as a white solid (200 mg).

Step B

In a manner analogous to that described in Example 6 (step B), reaction of the material from
15 step A above (200 mg) with 10% Pd/C (60 mg) under a hydrogen atmosphere afforded (2R-cyclopentylmethyl-4-hydroxy-3S-(4-methoxyphenylsulfonyl)amino)succinyl-piperidinamide as a white solid (250 mg).

Step C

In a manner analogous to that described in Example 7 (step A), reaction of the material from
20 step B above (250 mg) with O-benzylhydroxylamine hydrochloride (110 mg), N-methylmorpholine (0.15 mL) and TBTU (215 mg) afforded (4-benzyloxyamino-2R-cyclopentylmethyl-3S-(4-methoxyphenylsulfonyl)amino)succinyl-piperidinamide (265 mg) as a white solid. H¹-NMR (400 MHz, CDCl₃) 0.7-1.7 (m, 17 H); 3.4-3.6 (m, 5 H); 3.8 (dd, 1 H, J= 2.9 and 6.4 Hz); 3.85 (s, 3 H); 4.80 and 4.84 (two d, 2 H, J= 11.1 Hz); 6.95 (d, 2 H, J= 9.1 Hz);
25 7.19 (d, 1 H, J= 6.4 Hz); 7.39 (m, 5 H); 7.76 (d, 2 H, J= 9.1 Hz); 9.69 (s, 1 H) ppm. ESI/CID-MS 580 (MNa)⁺, 558 (MH)⁺, 435, 407, 388, 237, 149, 112 m/z.

Step D

In a manner analogous to that described in Example 7 (step B), reaction of the material from
step C above (260 mg) with 10% Pd/C (50 mg) under a hydrogen atmosphere afforded the
30 title compound (125 mg) as a white solid. H¹-NMR (200 MHz, CDCl₃) 0.8-1.8 (m, 17 H); 3.2-3.8 (m, 5 H); 3.86 (s, 3 H); 3.90 (m, 1 H); 6.98 (d, 2 H, J= 9.0 Hz); 7.02 (br signal, 1 H); 7.29 (s, 1 H); 7.83 (d, 2 H, J= 9.0 Hz); 9.81 (br signal, 1 H) ppm. FAB-MS 468 (MH)⁺, 435, 407, 298, 237, 208, 171, 112, 86, 84 m/z.

EXAMPLE 24**(3S-Amino-4-hydroxyamino-2R-isobutyl)succinyl-dicyclohexylmethanamide****Step A**

Sodium azide (66 mg) and dicyclohexylamine (300 mg; see Preparation G) were added at
5 room temperature to a solution of 4S-benzyloxycarbonyl-1-tert-butoxycarbonyl-3R-isobutyl-
azetidin-2-one (360 mg; see Preparation A) in DMF (8 mL). The mixture was stirred
overnight, then diluted with EtOAc and washed with H₂O, saturated aqueous NaHCO₃ and
2% HCl. The organic layer was dried over Na₂SO₄, filtered and evaporated. The crude
product was purified by flash chromatography on silica gel (n-hexane/EtOAc) to afford (4-
10 benzyloxy-3S-tert-butoxycarbonylamino-2R-isobutyl)succinyl-dicyclohexylmethanamide (230
mg) as a white solid. ESI-MS 567 (MNa)⁺, 545(MH)⁺, 489, 445, 167 m/z.

Step B

In a manner analogous to that described in Example 1 (step B), reaction of the material from
step A above (220 mg) with 10% Pd/C (50 mg) under a hydrogen atmosphere afforded (3S-
15 tert-butoxycarbonylamino-4-hydroxy-2R-isobutyl)succinyl-dicyclohexylmethanamide as an oil
(180 mg).

Step C

In a manner analogous to that described in Example 2 (step A), reaction of the material from
step B above (180 mg) with O-benzylhydroxylamine hydrochloride (77 mg), N-methyl-
20 morpholine (0.1 mL) and TBTU (128 mg) afforded (4-benzyloxyamino-3S-tert-butoxy-
carbonylamino-2R-isobutyl)succinyl-dicyclohexylmethanamide (200 mg) as a white solid. ¹H-
NMR (400 MHz, CDCl₃) 0.80-1.8 (m, 25 H); 0.92 (d, 6 H, J= 6.4 Hz); 1.39 (s, 9 H); 3.02
(m, 1 H); 3.56 (m, 1 H); 4.25 (d, 1 H, J= 7.9 Hz); 4.85 (s, 2 H); 5.58 (d, 1 H, J= 9.4 Hz);
6.41 (d, 1 H, J= 7.9 Hz); 7.34 (m, 5 H); 9.31 (s, 1 H) ppm. ESI-MS 610 (MK)⁺, 594
25 (MNa)⁺, 572 (MH)⁺, 516, 488, 472, 112 m/z.

Step D

In a manner analogous to that described in Example 2 (step B), reaction of the material from
step C above (180 mg) with 10% Pd/C (50 mg) under a hydrogen atmosphere afforded (3S-
tert-butoxycarbonylamino-4-hydroxyamino-2R-isobutyl)succinyl-dicyclohexylmethanamide
30 (100 mg) as a white solid.

Step E

In a manner analogous to that described in Example 3, reaction of the material from step D
above (100 mg) with 95% aqueous trifluoroacetic acid (5 mL) afforded the title compound,

trifluoroacetate salt, as a white solid (85 mg). In DMSO solution, the compound is present as a mixture of two rotamers (M, major; m, minor). ^1H -NMR (400 MHz, DMSO- d_6) 0.86 and 0.88 (two d, 6 H, J = 6.8 Hz); 0.9-1.7 (m, 25 H); 2.74 (m, 1 H, M); 3.10 (m, 1 H, m); 3.60 (m, 2 H); 7.76 (d, 1 H, J = 9.8 Hz); 8.01 (br signal, 3 H, M); 8.10 (br signal, 3 H, m); 9.25 (br
5 signal, 1 H, M); 9.57 (s, 1 H, m); 10.67 (s, 1 H, m); 11.00 (s, 1 H, M) ppm. FAB-MS 382 (MH) $^+$, 321 (M-HONHCO) $^+$, 194, 187, 112, 100, 89, 83 m/z.

EXAMPLE 25

(3S-Amino-4-hydroxyamino-2R-isobutyl)succinyl-diphenylmethanamide

Step A

10 In a manner analogous to that described in Example 28 (step A), reaction of 4S-benzyloxycarbonyl-1-tert-butoxycarbonyl-3R-isobutylazetidin-2-one (360 mg; see Preparation A) with diphenylmethanamine (0.34 ml) and NaN_3 (33 mg) afforded (4-benzyloxy-3S-tert-butoxycarbonylamino-2R-isobutyl)succinyl-diphenylmethanamide (150 mg) as a white solid. ^1H -NMR (400 MHz, CDCl_3) 0.89 and 0.91 (two d, 6 H, J = 6.7 Hz); 1.2-1.7 (m, 3 H); 1.39
15 (s, 9 H); 2.97 (m, 1 H); 4.46 (dd, 1 H, J = 3.8 and 9.7 Hz); 4.95 and 5.02 (two d, 2 H, J = 12.6 Hz); 6.07 (d, 1 H, J = 9.7 Hz); 6.09 and 6.14 (two d, 2 H, J = 7.6 Hz); 7.1-7.4 (m, 15 H) ppm.

Step B

In a manner analogous to that described in Example 1 (step B), reaction of the material from
20 step A above (150 mg) with 10% Pd/C (30 mg) under a hydrogen atmosphere afforded (3S-tert-butoxycarbonylamino-4-hydroxy-2R-isobutyl)succinyl-diphenylmethanamide as a white solid (100 mg).

Step C

In a manner analogous to that described in Example 2 (step A), reaction of the material from
25 step B above (100 mg) with O-benzylhydroxylamine hydrochloride (42 mg), N-methylmorpholine (0.06 mL) and TBTU (85 mg) afforded (4-benzyloxyamino-3S-tert-butoxycarbonylamino-2R-isobutyl)succinyl-diphenylmethanamide (100 mg) as a white solid.

Step D

In a manner analogous to that described in Example 2 (step B), reaction of the material from
30 step C above (100 mg) with 10% Pd/C (20 mg) under a hydrogen atmosphere afforded (3S-tert-butoxycarbonylamino-4-hydroxyamino-2R-isobutyl)succinyl-diphenylmethanamide as a white solid (90 mg).

Step E

In a manner analogous to that described in Example 3, reaction of the material from step D above (90 mg) with 95% aqueous trifluoroacetic acid (5 mL) afforded the title compound as a white solid (85 mg; trifluoroacetate salt). In DMSO solution, the compound is present as mixture of two rotamers (M, major; m, minor). H^1 - NMR (400 MHz, DMSO- d_6) 0.74 and 0.84 (two d, 6 H, J = 6.8 Hz); 1.12 and 1.49 (two m, 2 H); 1.31 (m, 1 H); 2.85 (m, 1 H); 3.58 (m, 1 H); 6.09 (d, 1 H, J = 7.7 Hz); 7.28 (m, 10 H); 8.20 (br signal, 3 H); 9.13 (d, 1 H, J = 8.1 Hz, M); 9.25 (d, 1 H, J = 8.1 Hz, m); 9.38 (br signal, 1 H, M); 9.60 (s, 1 H, m); 10.80 (s, 1 H, m); 11.10 (s, 1 H, M) ppm. FAB-MS 370 (MH)⁺, 309 (M-HONHCO)⁺, 204, 182, 167, 100 m/z.

10 EXAMPLE 26

(3S-tert-Butoxycarbonylamino-2R-cyclopentylmethyl-4-hydroxyamino)succinyl-(4-piperonyl)piperazinamide

Step A

In a manner analogous to that described in Example 1 (step A), reaction of 4S-benzyloxycarbonyl-1-tert-butoxycarbonyl-3R-cyclopentylmethylazetidin-2-one (500 mg; see Preparation D) with 1-piperonylpiperazine (545 mg) in acetonitrile afforded (4-benzyloxycarbonyl-3S-tert-butoxycarbonylamino-2R-cyclopentylmethyl)succinyl-(4-piperonyl)piperazinamide (800 mg) as an oil. H^1 - NMR (200 MHz, CDCl₃) 1.0-1.9 (m, 11 H); 1.41 (s, 9 H); 2.30 (m, 4 H); 3.36 (s, 2 H); 3.36 (m, 1 H); 3.50 (m, 4 H); 4.51 (dd, 1 H, J = 4.1 and 9.8 Hz); 5.06 and 5.18 (two d, 2 H, J = 12.4 Hz); 5.94 (s, 2 H); 6.40 (d, 1 H, J = 9.8 Hz); 6.6-6.9 (m, 3 H); 7.33 (m, 5 H) ppm. ESI/CID-MS 630 (MNa)⁺, 608 (MH)⁺, 508, 372, 221, 135 m/z.

Step B

In a manner analogous to that described in Example 1 (step B), reaction of the material from step A above (800 mg) with 10% Pd/C (100 mg) under a hydrogen atmosphere afforded (3S-tert-butoxycarbonylamino-2R-cyclopentylmethyl-4-hydroxy)succinyl-(4-piperonyl)piperazinamide (600 mg) as a white solid.

Step C

In a manner analogous to that described in Example 2 (step A), reaction of the material from step B above (600 mg) with O-benzylhydroxylamine hydrochloride (160 mg), N-methylmorpholine (0.27 mL) and TBTU (385 mg) afforded (4-benzyloxyaminocarbonyl-3S-tert-butoxycarbonylamino-2R-cyclopentylmethyl)succinyl-(4-piperonyl)piperazinamide (280 mg) as a white solid. H^1 - NMR (400 MHz, DMSO- d_6) 0.9-1.7 (m, 11 H); 1.37 (s, 9 H); 2.1-2.4

(m, 4 H); 3.18 (m, 1 H); 3.30-3.60 (m, 6 H); 4.01 (dd, 1 H, J= 6.4 and 9.0 Hz); 4.74 (s, 2 H); 5.97 (s, 2 H); 6.66 (d, 1 H, J= 9.0 Hz); 6.7-6.8 (m, 3 H); 7.34 (m, 5 H); 11.38 (s, 1 H) ppm. ESI-MS 645 (MNa)⁺, 623 (MH)⁺, 567, 523, 372, 283, 221 m/z.

Step D

- 5 In a manner analogous to that described in Example 2 (step B), reaction of the material from step C above (280 mg) with 10% Pd/C (50 mg) under a hydrogen atmosphere afforded the title compound (160 mg) as a white solid. H¹- NMR (400 MHz, DMSO-d₆) 0.9-1.7 (m, 11 H); 1.35 (s, 9 H); 2.1-2.4 (m, 4 H); 4.00 (dd, 1 H, J= 6.8 and 9.0 Hz); 5.97 (s, 2 H); 6.57 (d, 1 H, J= 9.0 Hz); 6.6-6.8 (m, 3 H); 8.80 (br signal, 1 H); 10.2 (br signal, 1 H) ppm. ESI-MS
10 555 (MNa)⁺, 533 (MH)⁺, 477, 444, 221 m/z.

EXAMPLE 27

(3S-Amino-2R-cyclopentylmethyl-4-hydroxyamino)succinyl-(4-piperonyl)piperazin- amide

- In a manner analogous to that described in Example 3, reaction of the compound described in
15 Example 26 (140 mg) with 95% aqueous trifluoroacetic acid (3 mL) afforded the title compound (bis-trifluoroacetate salt) as a white solid (100 mg). In DMSO solution, the compound is present as a mixture of two rotamers (M, major; m, minor). H¹- NMR (400 MHz, DMSO-d₆) 1.0-1.8 (m, 11 H); 2.8-4.6 (br m, 10 H); 3.20 (m, 1 H); 3.70 (m, 1 H); 6.07 (s, 2 H); 7.00 (m, 3 H); 8.08 (br signal, 3 H, m); 8.22 (br signal, 3 H, M); 9.43 (br
20 signal, 1 H, M); 10.1 (br signal, 1 H, M); 10.8 (s, 1 H, m); 11.20 (s, 1 H, M) ppm. ESI-MS 459 (MNa)⁺, 433 (MH)⁺, 299, 238, 221, 135 m/z.

By minor variations of the procedures of the examples above, and starting from the appropriate intermediates, the following compounds were obtained:

EXAMPLE 28

- 25 (3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-azetidinamide
Trifluoroacetate salt. H¹- NMR (400 MHz, DMSO-d₆): 1.44 (m, 4 H, CH₂CH₂-CH₂Ph), 2.14 (m, 2 H, azetidine 3-methylene), 2.47 (m, 2 H, CH₂CH₂-CH₂-Ph), 2.58 (m, 1 H, CH-phenpropyl), 3.57 (d, J= 6.1 Hz, 1 H, CH-NH₃⁺), 3.69 (s, 3 H, OMe), 3.86, 4.10 and 4.30 (each m; 4 H, azetidine 2-,4-methylene), 6.82 (d, J= 8.2 Hz, 2 H, aromatic protons ortho to methoxy), 7.07 (d, J= 8.2 Hz, 2 H, aromatic protons meta to methoxy), 8.00 (br s, 3 H, NH₃⁺), 9.32 (s, 1 H, NH-OH), and 11.10 ppm (br s, 1 H, NH-OH).

EXAMPLE 29

(3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-N-cyclohexylamide

Trifluoroacetate salt. H^1 - NMR (400 MHz, DMSO- d_6 ; 55 °C): 1.0-1.9 (m, 13 H, $\underline{CH_2CH_2-CH_2Ph}$ + cyclohexyl methylene protons), 2.50 (m, 3 H, $\underline{CH-CH_2CH_2-CH_2-Ph}$), 3.45 (d, $J=7.3$ Hz, 1 H, $\underline{CH-NH_3^+}$), 3.55 (m, 1 H, cyclohexyl methyne proton), 3.70 (s, 3 H, OMe), 6.81 (d, $J=8.6$ Hz, 2 H, aromatic protons ortho to methoxy), 7.04 (d, $J=8.6$ Hz, 2 H, aromatic protons meta to methoxy), 7.81 (d, $J=7.7$ Hz, 1 H, $\underline{CONH-cyclohexyl}$), 8.00 (br s, 3 H, $\underline{NH_3^+}$), 9.10 (br s, 1 H, $\underline{NH-OH}$), and 10.0 ppm (br s, 1 H, $\underline{NH-OH}$). At 55 °C, the $-NH_3^+$ protons were not detectable (7-8 ppm at lower temperatures).

EXAMPLE 30

(2R-Cyclopentylmethyl-3S-N,N-dimethylamino-4-hydroxyamino)succinyl-piperidin-
amide

H^1 - NMR (400 MHz, DMSO- d_6): 0.9-1.9 (m, 17 H, cyclopentylmethyl + piperidine 3-,4-,5-methylene protons), 2.12 (s, 6 H, $\underline{NMe_2}$), 3.00 (d, $J=10.7$ Hz, 1 H, $\underline{CH-NMe_2}$), 3.27 (m, 1 H, $\underline{CH-cyclopentylmethyl}$), 3.3-3.6 (m, 4 H, piperidine 2-,6-methylene), 8.83 (s, 1 H, $\underline{NH-OH}$), and 10.51 ppm (br s, 1 H, $\underline{NH-OH}$).

EXAMPLE 31

(4-Hydroxyamino-2R-(4-methoxy)phenpropyl-3S-phenylmethylsulfonylamino)succinyl-piperidinamide

H^1 - NMR (400 MHz, DMSO- d_6): 1.2-1.6 (m, 10 H, $\underline{CH-CH_2CH_2-CH_2Ph}$ + piperidine 3-,4-,5-methylene protons), 2.41 (m, 2 H, $\underline{CH_2-Ph}$), 3.16 (m, 1 H, $\underline{CH-phenpropyl}$), 3.44 (m, 4 H, piperidine 2-,6-methylene), 3.69 (s, 3 H, methoxy), 3.93 (m, 1 H, $\underline{CH-NHSO_2}$), 4.08 and 4.19 (each d, $J=13.7$ Hz, $\underline{NHSO_2-CH_2-Ph}$), 6.80 (d, $J=8.6$ Hz, 2 H, aromatic protons ortho to methoxy), 7.05 (d, $J=8.6$ Hz, 2 H, aromatic protons meta to methoxy), 7.33 (m, 5 H, Ph), 7.59 (br s, 1 H, $\underline{NH-SO_2}$), 9.02 (br s, 1 H, $\underline{NH-OH}$), and 10.90 ppm (br s, 1 H, $\underline{NH-OH}$).

EXAMPLE 32

(3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-(1R-cyclohexylethyl)-amide

Trifluoroacetate salt. H^1 - NMR (400 MHz, DMSO- d_6): 0.93 (d, $J=6.8$ Hz, 3 H, $\underline{CH_3-CH}$), 0.8-1.8 (m, 15 H, $\underline{CH-CH_2CH_2-CH_2Ph}$ + cyclohexyl protons), 2.43 (m, 2 H, $\underline{CH_2-Ph}$), 2.62 (m, 1 H, $\underline{CH-phenpropyl}$), 3.57 (m, 2 H, $\underline{CH-CH_3}$ + $\underline{CH-NH_3^+}$), 3.69 (s, 3 H, methoxy), 6.81 (d, $J=8.6$ Hz, 2 H, aromatic protons ortho to methoxy), 7.05 (d, $J=8.6$ Hz, 2 H, aromatic protons meta to methoxy), 7.93 (d, $J=8.6$ Hz, 1 H, $\underline{CO-NH-CH}$), 8.12 (br s, 3 H, $\underline{NH_3^+}$), 9.33 (s, 1 H, $\underline{NH-OH}$), and 11.05 ppm (s, 1 H, $\underline{NH-OH}$).

EXAMPLE 33

(3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-(1S-cyclohexylethyl)-amide

Trifluoroacetate salt. H^1 - NMR (400 MHz, DMSO- d_6): 0.8-1.8 (m, 15 H, $CH-CH_2CH_2-CH_2Ph$ + cyclohexyl protons), 0.97 (d, $J=6.8$ Hz, 3 H, CH_3-CH), 2.44 (m, 2 H, CH_2-Ph),
 5 2.61 (m, 1 H, CH -phenpropyl), 3.62 (m, 2 H, $CH-CH_3$ + $CH-NH_3^+$), 3.68 (s, 3 H, methoxy),
 6.80 (d, $J=8.5$ Hz, 2 H, aromatic protons ortho to methoxy), 7.03 (d, $J=8.5$ Hz, 2 H, aromatic protons meta to methoxy), 7.92 (d, $J=8.5$ Hz, 1 H, $CO-NH-CH$), 8.10 (br s, 3 H, NH_3^+), 9.34 (s, 1 H, $NH-OH$), and 11.08 ppm (s, 1 H, $NH-OH$).

EXAMPLE 34

10 **(3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-(4-tert-butylamino-carbonyl)piperidinamide**

Trifluoroacetate salt. In DMSO, the compound is present as two conformers, ca. 1:1, whose H^1 -NMR signals coalesce upon heating. H^1 -NMR (400 MHz, DMSO- d_6): 1.21 and 1.22 (each s, 9 H, t-Bu of 2 conformers), 1.2-1.8 (m, 8 H, $CH-CH_2CH_2-CH_2Ph$ + piperidine 3-,5-methylene), 2.30 (m, 1 H, piperidine 4-proton), 2.45 (m, 2 H, CH_2-Ph), 2.55 and 2.92 (each m; piperidine 2- and 6- axial protons), 3.20 (m, 1 H, CH -phenpropyl), 3.69 (s, 3 H, methoxy), 3.69 and 3.70 (each d, $J=7.0$ Hz; 1 H, $CH-NH_3^+$ of 2 conformers), 3.87 and 4.35 (each m; 2 H, piperidine 2- and 6- equatorial protons), 6.81 (d, $J=8.4$ Hz, 2 H, aromatic protons ortho to methoxy), 7.05 and 7.07 (each d, $J=8.4$ Hz; 2 H, aromatic protons meta to methoxy of 2 conformers), 7.35 (s, 1 H, $CO-NH-tBu$), 8.00 (br s, 3 H, NH_3^+), 9.34 and 9.37 (each s; 1 H, $NH-OH$ of 2 conformers), and 11.10 ppm (br s, 1 H, $NH-OH$).

EXAMPLE 35

(2R-Cyclopentylmethyl-3S-N,N-dimethylamino-4-hydroxyamino)succinyl-(4-piperonyl)-piperazinamide

25 H^1 - NMR (400 MHz, $CDCl_3$): 1.0-1.8 (m, 11 H, cyclopentylmethyl), 2.28 (s, 6 H, NMe_2), 2.38 and 2.46 (each m; 4 H, piperazinamide 3-,5-methylene), 3.35 (m, 2 H, CH -dimethylamino + CH -cyclopentylmethyl), 3.42 (s, 2 H, $N-CH_2-Ar$), 3.6-3.8 (m, 4 H, piperazinamide 2-,6-methylene), 5.95 (s, 2 H, $O-CH_2-O$), 6.75 and 6.85 (each m; 3 H, Ar), 8.00 (br s, 1 H, $NH-OH$), and 10.00 ppm (br s, 1 H, $NH-OH$).

30 **EXAMPLE 36**

(3S-(N-Cyclohexylmethyl-N-methyl)amino-4-hydroxyamino-2R-(4-methoxy)-phenpropyl)-succinyl-piperidinamide

H^1 - NMR (400 MHz, DMSO- d_6): 0.6-1.8 (m, 21 H, CH-CH₂CH₂-CH₂Ph + piperidine 3-,4-,5-methylene + cyclohexyl protons), 2.10 (m, 2 H, CH₂-Ph), 2.12 (s, 3 H, NMe), 2.40 (m, 2 H, N-CH₂-cyclohexyl), 3.06 (d, J= 10.7 Hz, 1 H, CO-CH-N), 3.2-3.5 (m, 5 H, CH-phenpropyl + piperidine 2-,6-methylene), 3.69 (s, 3 H, methoxy), 6.80 (d, J= 8.5 Hz, 2 H, aromatic protons ortho to methoxy), 7.02 (d, J= 8.5 Hz, 2 H, aromatic protons meta to methoxy), and 9.00 ppm (br s, 1 H, NH-OH); NH-OH not detected.

EXAMPLE 37

(2R-Cyclopentylmethyl-4-hydroxyamino-3S-(4-methoxyphenyl)sulfonylamino)succinyl-(4-piperonyl)-piperazinamide

Trifluoroacetate salt. In DMSO, the compound is present as a mixture of two conformers (coalescence of NMR signals upon heating). H^1 - NMR (400 MHz, DMSO- d_6): 0.8-1.6 (m, 11 H, cyclopentylmethyl), 2.5-3.5 (m, 7 H, CH-cyclopentylmethyl + piperazinamide 3-,5-methylene + CH₂-Ar), 3.5-4.5 (m, 5 H, CH-NH SO_2 + piperazinamide 2-,6-methylene), 6.07 (s, 2 H, O-CH₂-O), 6.9-7.1 (m, 5 H, piperonyl Ar protons + aromatic protons ortho to methoxy), 7.63 (d, J= 8.6 Hz, 2 H, aromatic protons meta to methoxy), 7.73 and 7.86 (d, J= 9.0 Hz, NH- SO_2), 8.89 (br s, 1 H, NH^+), 9.70 and 9.80 (br s, 1 H, NH-OH), and 10.77, 10.82 ppm (br s, 1 H, NH-OH).

EXAMPLE 38

2R-(4S-(2-Dimethyl-1-hydroxy-2-oxo-3H-imidazolidinyl)-3-cyclopentylpropionic acid (4-piperonyl)-piperazinamide

Obtained as a white solid by stirring (3S-amino-2R-cyclopentylmethyl-4-hydroxyamino)-succinyl-(4-piperonyl)piperazinamide (Example 27) in neat acetone, followed by removal of water and solvent by rotoevaporation with anhydrous ethanol. Trifluoroacetate salt. H^1 - NMR (400 MHz, DMSO- d_6): 1.0-1.8 (m, 11 H, cyclopentylmethyl), 1.16 and 1.17 (each s, 3 H, Me), 2.9-4.4 (m, 10 H, CH-cyclopentylmethyl + piperazine methylene protons + CH₂-Ar + CH-NH $_2^+$), 6.05 (s, 2 H, O-CH₂-O), 6.9-7.1 (m, 3 H, piperonyl Ar protons), 9.64 (s, 1 H, N-OH), and 9.9 ppm (br s, 2 H, -NH $_2^+$).

EXAMPLE 39

(4-Hydroxyamino-2R-(4-methoxy)phenpropyl-3S-(3-pyridyl)methylamino)succinyl-piperidinamide

H^1 - NMR (400 MHz, DMSO- d_6): 1.1-1.6 (m, 10 H, CH-CH₂CH₂-CH₂Ph + piperidine 3-,4-,5-methylene protons), 2.40 (m, 2 H, CH₂-Ph), 3.06 (m, 2 H, CH-NH + CH-phenpropyl), 3.2-3.6 (m, 4 H, piperidine 2-,6-methylene), 3.37 and 3.70 (each d, J= 16.2 Hz; 2 H, NH-

CH₂-pyridine), 3.68 (s, 3 H, methoxy), 6.79 (d, J= 7.6 Hz, 2 H, aromatic protons ortho to methoxy), 7.10 (d, J= 7.6 Hz, 2 H, aromatic protons meta to methoxy), 7.26 (dd, J= 4.7 and 7.7 Hz, 1 H, 5-pyridine proton), 7.60 (ddd, J= 1.7, 2.1 and 7.7 Hz, 1 H, 4-pyridine proton), 8.38 (dd, J= 1.7 and 4.7 Hz, 1 H, 6-pyridine proton), and 8.39 ppm (d, J= 2.1 Hz, 1 H, 2-pyridine proton); NH-OH and NH-OH not detected.

EXAMPLE 40**N-((3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl)-L-proline-piperonylamide**

Trifluoroacetate salt. ¹H- NMR (400 MHz, DMSO-d₆): 1.4-2.1 (m, 8 H, CH-CH₂CH₂-CH₂Ph + proline 3-,4-methylene), 2.45 (m, 2 H, CH₂-Ph), 2.95 (m, 1 H, CH-phenpropyl), 3.5-3.7 (m, 3 H, CH-NH₃⁺ + proline 5-methylene), 3.67 (s, 3 H, methoxy), 4.16 and 4.21 (each dd, J= 5.6 and 13.7 Hz, 1 H, CONH-CH(H)-Ar), 4.34 (dd, J= 4.3 and 8.1 Hz, proline N-CH-CO), 6.07 (s, 2 H, O-CH₂-O), 6.6-6.9 (m, 5 H, piperonyl Ar protons + aromatic protons ortho to methoxy), 7.06 (d, J= 8.6 Hz, 2 H, aromatic protons meta to methoxy), 8.15 (br s, 3 H, NH₃⁺), 8.31 (dd, J= 5.6 Hz, 1 H, CO-NH-CH₂-Ar), 9.39 (s, 1 H, NH-OH), and 11.16 ppm (s, 1 H, NH-OH).

EXAMPLE 41**(3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-N-isopropyl-N-methylamide, and****(3R-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-N-isopropyl-N-methylamide**

A solution of 4S-benzyloxycarbonyl-1-tert-butoxycarbonyl-3R-(4-methoxy)phenpropyl-azetidin-2-one (350 mg; see Preparation C) in DMF (6 mL) was allowed to react with sodium azide (10 mg) and N-isopropyl-N-methylamine (0.16 mL) for 30 hr at room temperature. Workup and flash chromatography over silica (n-hexane/EtOAc) afforded, in this order, (4-benzyloxycarbonyl-3S-tert-butoxycarbonylamino-2R-(4-methoxy) phenpropyl) succinyl-N-isopropyl-N-methylamide (160 mg) and the product from base-assisted epimerization, (4-benzyloxycarbonyl-3R-tert-butoxycarbonylamino-2R-(4-methoxy) phenpropyl) succinyl-N-isopropyl-N-methylamide (80 mg). ESI-MS (both isomers) 527 (MH)⁺.

Separate elaboration of the two diastereoisomers above, in a manner analogous to that described in Examples 1-3, afforded the title compounds (both trifluoroacetate salts). In

DMSO, both compounds exist as a 2:1 mixture (M, major; m, minor; H^1 - NMR) of rotamers at the tertiary amide.

3S-Isomer: H^1 - NMR (600 MHz, DMSO- d_6): 0.97 and 1.02 (each d, $J = 7.0$ Hz; 6 H, $CHMe_2$ of M), 1.01 and 1.11 (each d, $J = 7.0$ Hz; 6 H, $CHMe_2$ of m), 1.47 (m, 4 H, $CH-CH_2CH_2-CH_2Ph$), 2.44 (m, 2 H, CH_2-Ph), 2.65 (s, 3 H, NMe of m), 2.77 (s, 3 H, NMe of M), 3.11 (m, 1 H, CH -phenpropyl of M), 3.18 (m, 1 H, CH -phenpropyl of m), 3.69 (s, 3 H, methoxy), 3.71 (m, 1 H, $CH-NH_3^+$ of M), 3.76 (m, 1 H, $CH-NH_3^+$ of m), 4.03 (m, 1 H, $CH-Me_2$ of m), 4.69 (m, 1 H, $CH-Me_2$ of M), 6.82 (d, $J = 8.6$ Hz, 2 H, aromatic protons ortho to methoxy), 7.05 (d, $J = 8.6$ Hz, 2 H, aromatic protons meta to methoxy), 7.98 (br s, 3 H, NH_3^+ of m), 8.13 (br s, 3 H, NH_3^+ of M), 9.36 (s, 1 H, $NH-OH$ of M), 9.37 (s, 1 H, $NH-OH$ of m), 11.12 (s, 1 H, $NH-OH$ of M), and 11.15 ppm (s, 1 H, $NH-OH$ of m).

3R-Isomer: H^1 - NMR (600 MHz, DMSO- d_6): 0.98 and 1.00 (each d, $J = 6.5$ Hz; 6 H, $CHMe_2$ of M), 1.04 and 1.11 (each d, $J = 6.5$ Hz; 6 H, $CHMe_2$ of m), 1.3-1.8 (m, 4 H, $CH-CH_2CH_2-CH_2Ph$), 2.46 (m, 2 H, CH_2-Ph), 2.64 (s, 3 H, NMe of m), 2.80 (s, 3 H, NMe of M), 3.17 (m, 1 H, CH -phenpropyl), 3.59 (m, 1 H, $CH-NH_3^+$), 3.69 (s, 3 H, methoxy of M), 3.70 (s, 3 H, methoxy of m), 4.12 (m, 1 H, $CH-Me_2$ of m), 4.67 (m, 1 H, $CH-Me_2$ of M), 6.81 (d, $J = 8.5$ Hz, 2 H, aromatic protons ortho to methoxy of M), 6.83 (d, $J = 8.5$ Hz, 2 H, aromatic protons ortho to methoxy of m), 7.05 (d, $J = 8.5$ Hz, 2 H, aromatic protons meta to methoxy of M), 7.09 (d, $J = 8.5$ Hz, 2 H, aromatic protons meta to methoxy of m), 8.21 (br s, 3 H, NH_3^+), 9.20 (br s, 1 H, $NH-OH$ of m), 9.22 (br s, 1 H, $NH-OH$ of M), and 10.99 ppm (br s, 1 H, $NH-OH$).

EXAMPLE 42

(3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-4-(2-pyridyl)-piperazinamide

Trifluoroacetate salt. H^1 - NMR (400 MHz, DMSO- d_6): 1.48 (m, 4 H, $CH-CH_2CH_2-CH_2Ph$), 2.44 (m, 2 H, CH_2-Ph), 3.22 (m, 1 H, CH -phenpropyl), 3.3-3.8 (m, 9 H, $CH-NH_3^+$ and piperidine methylene protons), 3.68 (s, 3 H, methoxy), 6.70 (dd, $J = 5.1$ and 6.8 Hz, 5-pyridine proton), 6.78 (d, $J = 8.6$ Hz, 2 H, aromatic protons ortho to methoxy), 6.88 (d, $J = 8.6$ Hz, 1 H, 3-pyridine proton), 7.05 (d, $J = 8.6$ Hz, 2 H, aromatic protons meta to methoxy), 7.60 (ddd, $J = 1.7$, 6.8 and 8.6 Hz, 1 H, 4-pyridine proton), 8.10 (dd, $J = 1.7$ and 5.1 Hz, 1 H, 6-pyridine proton), 8.20 (br s, 3 H, NH_3^+), 9.40 (br s, 1 H, $NH-OH$), and 11.19 ppm (s, 1 H, $NH-OH$).

EXAMPLE 43

(3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-(4-(2,3-methylene-dioxy)phenylaminocarbonyl)piperidinamide

Trifluoroacetate salt. H^1 -NMR (400 MHz, DMSO- d_6): In DMSO, the compound is present as two conformers, ca. 1:1, whose H^1 -NMR signals coalesce upon heating. H^1 -NMR (400 MHz, DMSO- d_6): 1.2-1.9 (m, 8 H, $CH-CH_2CH_2-CH_2Ph$ + piperidine 3-,5-methylene), 2.45 (m, 2 H, CH_2-Ph), 2.50 (m, 1 H, piperidine 4-proton), 2.70 and 3.00 (each m; piperidine 2- and 6- axial protons), 3.20 (m, 1 H, CH -phenpropyl), 3.66 and 3.70 (each s; 3 H, methoxy of 2 conformers), 3.70 (m, 1 H, $CH-NH_3^+$), 3.90 and 4.40 (each m; 2 H, piperidine 2- and 6-equatorial protons), 5.95 (s, 2 H, $O-CH_2-O$), 6.8-7.3 (m, 7 H, aromatic protons), 8.00 (br s, 3 H, NH_3^+), 9.38 and 9.39 (each s; 1 H, $NH-OH$ of 2 conformers), 9.82 and 9.83 (each s; 1 H, $CO-NH-Ar$), and 11.15 ppm (br s, 1 H, $NH-OH$).

In a manner analogous to that described in the previous Examples, there were prepared the following compounds:

EXAMPLE 44

(3S-Amino-4-hydroxyamino-2R-isobutyl)succinyl-N-methyl-N- (1S-methylaminocarbonyl-2-phenyl)ethylamide, trifluoroacetate salt.

H^1 -NMR (600 MHz, DMSO- d_6): 0.73, 0.80 (two d, $J=6.3$ Hz, 6H), 1.1-1.5 (m, 3H), 2.55 (d, $J=4.4$ Hz, 3H), 2.83 (s, 3H), 3.05, 3.30 (two m, 2H), 3.10 (m, 1H), 3.60 (m, 1H), 4.70 (m, 1H), 7.0-7.3 (m, 5H), 7.81 (q, $J=4.4$ Hz, 1H), 8.28 (broad signal, 3H), 9.40 (broad signal, 1H), 11.13 (s, 1H).

EXAMPLE 45

(3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-(4-piperonyl)piperazinamide

EXAMPLE 46

(3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-4-(4-fluorophenylsulfonyl) piperazinamide

EXAMPLE 47

(2R-Cyclopentylmethyl-4-hydroxyamino-3S-(methylsulfonyl)amino)succinyl-(4-piperonyl)piperazinamide

EXAMPLE 48

(3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-4-(4-acetylphenyl)piperazinamide

EXAMPLE 49

(3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-4-(1-piperidyl)piperidinamide

EXAMPLE 50

5 (3S-Amino-2R-cyclopentylmethyl-4-hydroxyamino)succinyl-(1R-cyclohexylethyl)amide, trifluoroacetate salt.

^1H -NMR (400 MHz, DMSO- d_6): 0.9-1.9 (m, 22H); 0.95 (d, J= 6.8 Hz, 3H), 2.60 (m, 1H), 3.55 (m, 2H), 7.87 (d, J=7.1Hz, 1H), 8.0 (broad signal, 3H) 9.27 (s, 1H), 11.01 (broad signal, 1H).

EXAMPLE 51

10 (4-Hydroxyamino-3S-(4-methoxyphenylsulfonyl)amino-2R-(4-methoxy)phenpropyl)succinyl-piperidinamide

EXAMPLE 52

(3S-(4-Fluorophenylsulfonyl)amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-piperidinamide

15 **EXAMPLE 53**

(3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-cyclohexylamide

EXAMPLE 54

(3S-Amino-2R-cyclopentylmethyl-4-hydroxyamino)succinyl-piperazinamide

EXAMPLE 55

20 (3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-cyclopentylamide

EXAMPLE 56

(3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-(2R,S-methyl)piperidinamide

EXAMPLE 57

25 (3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-2-(1-morpholino)ethylamide

EXAMPLE 58

(3S-Amino-2R-cyclopentylmethyl-4-hydroxyamino)succinyl-2-(N-methyl-N-piperonyl)aminoethylamide

30 **EXAMPLE 59**

(3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-4-(1,1-di(4-fluoro)phenylmethyl)piperazinamide

EXAMPLE 60

35 (3S-Amino-2R-cyclopentylmethyl-4-hydroxyamino)succinyl-N-methyl-N-(2-(N-methyl-N-piperonyl)-aminoethylamide

EXAMPLE 61

(2R-Cyclopentylmethyl-3S-(4-fluorophenylsulfonyl)amino-4-hydroxyamino)succinyl-piperidinamide

EXAMPLE 62

- 5 **(3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-4-(4-piperidinecarbonyl)piperazinamide**

EXAMPLE 63

(3S-Amino-2R-cyclopentylmethyl-4-hydroxyamino)succinyl-(3,5-dimethyl)piperazinamide

- 10 **EXAMPLE 64**

(3S-Amino-2R-biphenpropyl-4-hydroxyamino)succinyl-(4-piperonyl)piperazinamide, trifluoroacetate salt.

- ¹H-NMR (500 MHz, DMSO-d₆): 1.60 (m, 4H), 2.60, 2.70 (two m, 2H), 3.0-4.0 (broad signal, 10H), 3.30 (m, 1H), 4.02 (d, J=7.2 Hz, 1H), 5.93 (s, 2H), 6.6-6.8 (m, 3H), 7.30-7.70 (m, 9H).

EXAMPLE 65

(3S-Amino-2R-cyclopentylmethyl-4-hydroxyamino)succinyl-(1R-cyclohexyl-2-hydroxy)ethylamide

EXAMPLE 66

- 20 **(3S-Amino-4-hydroxyamino-2R-(4-methoxy)phenpropyl)succinyl-4-(2-hydroxyethyl)piperazinamide**

EXAMPLE 67

N-((3S-Amino-2R-cyclopentylmethyl-4-hydroxyamino)succinyl)-L-proline--N-(2,3-methylenedioxyphenyl)-carboxamide

- 25 **EXAMPLE 68**

N-((3S-Amino-2R-cyclopentylmethyl-4-hydroxyamino)succinyl)-D-proline--N-(2,3-methylenedioxyphenyl)-carboxamide

EXAMPLE 69

- 30 **(2R-Cyclopentylmethyl-4-hydroxyamino-3S-(toluenesulfonyl)amino)succinyl-(1R-cyclohexylethyl)amide**

EXAMPLE 70

(3S-Amino-2R-cyclopentylmethyl-4-hydroxyamino)succinyl-(2S-methyl-4-piperonyl)piperazinamide

EXAMPLE 71

(2R-Cyclopentylmethyl-4-hydroxyamino-3S-(8-(1,2,3,4-tetrahydroquinoline)sulfonyl)amino)succinyl-(1R-cyclohexylethyl)amide

H^1 - NMR (400 MHz, DMSO- d_6): 0.89 (d, J = 6.8 Hz, 3H), 0.9-1.9 (m, 22H), 2.43 (m, 1H), 2.67 (m, 2H), 3.30 (m, 2H), 3.50 (m, 2H), 5.94 (s, 1H), 6.45 (dd J = 7.3, 7.8 Hz, 1H), 7.03 (d, J = 7.3 Hz, 1H), 7.30 (d, J =7.8 Hz, 1H), 7.3, 7.6 (two broad signals, 2H), 8.72 (broad signal, 1H) 10.50 (broad signal, 1H).

EXAMPLE 72

(3S-Amino-2R-cyclopentylmethyl-4-hydroxyamino)succinyl- (1S-cyclohexyl-1-dimethylaminocarbonyl)-methanamide

H^1 - NMR (400 MHz, DMSO- d_6): 0.9-1.9 (m,22H), 2.68 (m, 1H), 2.79,3.06 (two singlets, 6H), 3.47 (m, 1H), 4.50 (dd, J = 7.3, 8.8 Hz, 1H), 7.80 (broad signal, 3H), 8.16 (d, J =7.3 Hz, 1H), 9.21 (broad signal, 1H), 11.00 (broad signal, 1H).

EXAMPLE 73

(3S-Amino-2R-biphenpropyl-4-hydroxyamino)succinyl-(1R-cyclohexylethyl)amide, trifluoroacetate salt.

H^1 - NMR (400 MHz, DMSO- d_6): 0.90-1.9 (m, 15H), 0.94 (d, J = 6.8 Hz, 3H), 2.57 (m, 2H), 2.62 (m, 1H), 3.57 (m, 1H), 3.67 (m, 1H), 7.25, 7.55 (two doublets, J =8.3 Hz, 4H), 7.30 (m, 1H), 7.42 (m, 2H), 7.60 (m, 2H), 8.02 (d, J =8.8 Hz, 1H), 8.18 (broad signal, 3H), 9.32 (s, 1H), 11.20 (s, 1H).

PREPARATION A

4S-Benzyloxycarbonyl-1-tert-butoxycarbonyl-3R-isobutylazetidin-2-one

Step A

A solution of 1-tert-butyldimethylsilyl-4S-carboxyazetidin-2-one (6.2 g) in dry THF (100 mL) was treated dropwise at 0-5°C with a 2M solution of LDA (28.4 ml) in the same solvent, to obtain an orange solution of the di-anion. After 15 min, isobutyl iodide (6.8 mL) was added at 0°C under stirring, and the resulting yellow-green solution was left at the same temperature overnight. Quenching with 1M aqueous $KHSO_4$ (300 mL), followed by extraction with EtOAc, afforded crude 1-tert butyldimethylsilyl-4S-carboxy-3R-isobutylazetidin-2-one as an orange syrup (7 g).

The above material was dissolved in dry DMF (20 mL) and treated dropwise, in this order, with triethylamine (5.85 mL) and benzyl bromide (4.8 mL). After 4 hours at room temperature, the mixture was partitioned between water and EtOAc. The organic layer was washed with brine, dried and evaporated to give crude 4S-benzyloxycarbonyl-1-tert-butyldimethylsilyl-3R-isobutylazetidin-2-one as an orange oil. This material was dissolved in

THF (10 mL) and left overnight in the presence of tetrabutylammonium fluoride (2.6 g) and acetic acid (1.7 mL). The mixture was partitioned between saturated aqueous NaHCO₃ and EtOAc, and the organic phase was dried and evaporated. Flash chromatography over silica gel (n-hexane/EtOAc) afforded 4S-benzyloxycarbonyl-3R-isobutylazetidin-2-one (4.7 g) as white needles. FT-IR (KBr) 3229 (N), 1744-1750 br (CO) cm⁻¹. H¹-NMR (200 MHz, CDCl₃) 0.87 (d, 3 H, J= 6.5 Hz); 0.94 (d, 3 H, J= 6.5 Hz); 1.57-1.82 (m, 3 H); 3.32 (m, 1 H); 3.90 (d, 1 H, J= 2.4 Hz); 5.22 (ABq, 2 H); 5.96 (br s, 1 H); 7.36 (m, 5 H) ppm.

Step B

A solution of the material from step A above (1 g) in CH₃CN (15 mL) was treated with DMAP (4-dimethylaminopyridine; 46 mg) and Boc₂O (di-tert-butylidicarbonate; 1.67 g) at 40 °C for 30 min and then at room temperature overnight. After removal of the solvent in vacuo, the residue was dissolved in EtOAc and sequentially washed with aqueous 1M KHSO₄, aqueous saturated NaHCO₃ and brine. Drying over Na₂SO₄ and evaporation left the title compound as a yellow oil. FT-IR (CHCl₃) 1820 (azetidinone CO), 1750 (ester CO); 1728 (carbamate CO) cm⁻¹.

PREPARATION B

3R-Isobutyl-4S-(4-nitro)benzyloxycarbonyl-1-(4-toluenesulfonyl)azetidin-2-one

Step A

In a manner analogous to that described in Preparation A, reaction of 1-tert-butyl-dimethylsilyl-4S-carboxyazetidin-2-one (4.1 g) with a 2M solution of LDA in THF (18.8 mL) and isobutyl iodide (4.53 mL), followed by reaction with p-nitrobenzyl bromide (4.25 g) and triethylamine (3.7 mL), and finally by reaction with acetic acid (1 mL) and tetrabutylammonium fluoride (1.1 g), afforded 3R-isobutyl-4S-(4-nitro)benzyloxycarbonyl-azetidin-2-one (3.6 g) as a yellow solid.

Step B

A solution of the material from Step A above (630 mg) in CH₂Cl₂ (10 mL) was treated with triethylamine (0.65 ml) and p-toluenesulfonyl chloride (783 mg) at room temperature overnight under a nitrogen atmosphere. After quenching with saturated aqueous NaHCO₃, the organic layer was collected, washed with aqueous saturated NH₄Cl, brine and dried over Na₂SO₄. Evaporation and purification by flash chromatography on silica gel (n-hexane/EtOAc) afforded the title compound (320 mg) as an oil. FT-IR (CHCl₃) 1802 (azetidinone CO), 1752 (ester CO) cm⁻¹. H¹-NMR (400 MHz, CDCl₃) 0.79 (d, 3 H, J= 6.4 Hz); 0.88 (d, 3 H, J= 6.4

Hz); 1.54-1.72 (m, 3 H); 2.44 (s, 3 H); 3.20 (m, 1 H); 4.32 (d, 1 H, J= 3.2 Hz); 5.19 (s, 2 H); 7.31 (d, 2 H, J= 8.5 Hz); 7.33 (m, 5 H), 7.87 (d, 2 H, J= 8.5) ppm.

PREPARATION C

4S-Benzyloxycarbonyl-1-tert-butoxycarbonyl-3R-(4-methoxy)phenpropylazetidin-2-one

Step A

In a manner analogous to that described in Preparation A, reaction of 1-tert-butyl-dimethylsilyl-4S-carboxyazetidin-2-one (21 g) with a 2M solution of LDA in THF (93.8 mL) and (4-methoxy)phenpropyl iodide (45.9 g; see Preparation F), followed by reaction with benzyl bromide (32.6 mL) and triethylamine (15.3 mL), afforded crude 4S-benzyloxy-carbonyl-1-tert-butyl-dimethylsilyl-3R-(4-methoxy)phenpropylazetidin-2-one (90.4 g).

This material, dissolved in CH₃CN and aqueous 2N HCl, was let stirring at room temperature overnight. The solvent was evaporated and the aqueous phase was extracted with EtOAc several times. The organic layer was dried over Na₂SO₄ and evaporated. The residue was purified by flash chromatography on silica gel (n-hexane/EtOAc) to afford 4S-benzyloxycarbonyl-3R-(4-methoxy)phenpropylazetidin-2-one (8.6 g) as a white solid.

Step B

In a manner analogous to that described in Preparation A (step B), reaction of the material from step A above (8.6 g) with DMAP (300 mg) and Boc₂O (7.93 g) afforded the title compound as a yellow oil (9.1 g). ¹H-NMR (200 MHz, CDCl₃) 1.44 (s, 9 H); 1.6-1.9 (m, 4 H); 2.55 (m, 2 H); 3.15 (m, 1 H); 3.79 (s, 3 H); 4.09 (d, 1 H, J=3.2 Hz); 5.19 and 5.25 (two d, 2 H, J= 12.2 Hz); 6.81 (d, 2 H, J= 8.7 Hz); 7.02 (d, 2 H, J= 8.7 Hz); 7.34 (m, 5 H) ppm.

PREPARATION D

4S-Benzyloxycarbonyl-1-tert-butoxycarbonyl-3R-cyclopentylmethylazetidin-2-one

Step A

In a manner analogous to that described in Preparation A, reaction of 1-tert-butyl-dimethylsilyl-4S-carboxyazetidin-2-one (1 g) with a 2M solution of LDA in THF (4.6 mL) and cyclopentymethyl iodide (2 g; see Preparation E), followed by treatment with benzyl bromide (0.93 mL) and triethylamine (1 mL), afforded crude 4S-benzyloxycarbonyl-1-tert-butyl-dimethylsilyl-3R-cyclopentylmethylazetidin-2-one (6 g).

Step B

This material, dissolved in CH₃CN and aqueous 2N HCl, was let stirring at room temperature overnight. The solvent was evaporated and the aqueous phase was extracted with EtOAc

several times. The organic layer was dried over Na_2SO_4 and evaporated. The residue was purified by flash chromatography on silica gel (n-hexane/EtOAc) to afford 4S-benzyloxy-carbonyl-3R-cyclopentylmethylazetidin-2-one (1.3 g) as a yellow solid.

Step C

- 5 In a manner analogous to that described in Preparation A (step B), reaction of the material from step B above (1.3 g) with DMAP (21 mg) and Boc_2O (566 mg) afforded the title compound as a colourless oil (530 mg). ^1H -NMR (200 MHz, CDCl_3) 0.9-2.0 (m, 11 H); 1.45 (s, 9 H); 3.15 (m, 1 H); 4.12 (d, 1 H, $J = 3.0$ Hz); 5.17 and 5.30 (two d, 2 H, $J = 12.0$ Hz); 7.35 (m, 5 H) ppm. FAB- MS 388 (MH)⁺, 332, 288, 252, 135, 91, 57 m/z.

10 PREPARATION E

Cyclopentylmethyl iodide

- Methanesulfonyl chloride (25.5 mL) was added dropwise to a solution of cyclopentane-methanol (32.4 mL) and triethylamine (46 mL) in dichloromethane (500 mL) at 0 °C. After stirring overnight at room temperature, the mixture was sequentially washed with water, 2%
15 hydrochloric acid, 4% aqueous sodium bicarbonate and water again. Following drying over sodium sulfate, the solvent was removed under vacuum, affording crude cyclopentylmethyl methanesulfonate (ca. 50 g) as a colourless oil. A mixture of this product and sodium iodide in acetone (600 mL) was heated at reflux in the dark for 16 hours. After cooling to room temperature, the reaction mixture was poured into petrol ether/water 2:1 (1.5 L) under
20 stirring. The organic layer was washed twice with water, dried (Na_2SO_4) and carefully concentrated under reduced pressure (50 mm; temperature of the bath <30 °C). The residue was distilled in vacuo. The title product was obtained as a light yellow oil (39 g), by collecting the fraction boiling at 101-102°/70mm.

PREPARATION F

25 3-(4-Methoxyphenyl)-1-propyl iodide

- Methanesulfonyl chloride (31 mL) was added dropwise to a solution of 3-(4-methoxyphenyl)-1-propanol (50 g) and triethylamine (56 mL) in dichloromethane (800 mL) at 0 °C. The resulting mixture was stirred overnight at room temperature. After washing with water, 2 % hydrochloric acid and 4 % aqueous sodium bicarbonate, the organic solution was
30 dried over sodium sulfate and rotoevaporated to yield crude 3-(4-methoxyphenyl)-1-propyl methanesulfonate as a colourless solid (ca. 83 g). This product and sodium iodide (75 g) in acetone (800 mL) were heated at reflux in the dark for 20 hours. After cooling to room temperature, the reaction mixture was poured into n-hexane/EtOAc/water 1:1:1 (1.5 L)

under stirring. The organic layer was washed with water, diluted aqueous sodium metabisulfite and 4 % aqueous sodium bicarbonate, then dried (Na_2SO_4) and concentrated under reduced pressure. The residue was purified by passing it through a short pad of silica gel (eluting with n-hexane/EtOAc 95:5). The title compound was obtained as a pale yellow solid (75 g).

PREPARATION G

Dicyclohexylmethyl amine

A solution of dicyclohexylacetic acid (4.5 g) in dichloromethane (50 mL) was treated with oxalyl chloride (2.5 mL). Addition of a catalytic amount of N,N-dimethylformamide (1 drop) caused immediate gas evolution to take place. The reaction mixture was stirred for 2 hr, then concentrated to dryness. The residue, consisting of crude dicyclohexylacetyl chloride, was dissolved in acetone (45 mL) and treated at 0 °C with a solution of sodium azide (2.6 g) in water (45 mL). After stirring for 1 hr at 0 °C, toluene (100 mL) was added. The organic layer (mainly containing dicyclohexylacetyl azide as shown by IR absorption band at 2128 cm^{-1}) was separated, washed twice with brine, dried (Na_2SO_4), then heated at reflux for 2 hr. Following evaporation of the solvent, a waxy solid was obtained which mainly consisted of dicyclohexylmethyl isocyanate (IR absorption band at 2275 cm^{-1}). This crude isocyanate was suspended in 20% hydrochloric acid (100 mL) and heated at reflux for 8 h. The insoluble material was filtered off and the filtrate was concentrated in vacuo. The solid residue was taken up in diethyl ether /1N aqueous NaOH and the aqueous layer was re-extracted with ethyl ether. The organic phases were joined and washed with brine, then dried (Na_2SO_4) and rotoevaporated. The title product was thus obtained as a colourless oil (1.8 g). $^1\text{H-NMR}$ (200 MHz, DMSO-d_6) 0.8-1.8 (m, 22 H); 2.01 (t, 1 H, $J = 5.4$ Hz) ppm.

PREPARATION H

N-(tert-Butoxycarbonyl)-3R-isobutyl-L-aspartic acid 4-piperidinamide ("ASPARTIC ROUTE")

Step A

A solution of N-(tert-butoxycarbonyl)-L-aspartic acid 1-benzyl ester (1 g) and piperidine (0.32 mL) in acetonitrile (15 mL) was allowed to react with N-methylmorpholine (0.42 mL) and TBTU (1.2 g) for 6 hr at room temperature. The solvent was removed and the residue, dissolved in EtOAc, was sequentially washed with 2% aqueous hydrochloric acid, saturated aq. NaHCO_3 and brine. After drying (Na_2SO_4) and rotoevaporation, the resulting yellowish

oil was purified by flash chromatography over silica to obtain N-(tert-butoxycarbonyl)-L-aspartic acid 1-benzyl ester-4-piperidinamide (900 mg) as a colorless oil.

Step B

A solution of the compound from step A above (700 mg) in ethanol (30 mL) was exposed to a hydrogen atmosphere in the presence of 10% Pd-C (70 mg) for 4 hr at room temperature. After filtration (Celite filter aid) and rotoevaporation, N-(tert-butoxycarbonyl)-L-aspartic acid 4-piperidinamide (500 mg) was obtained as a yellowish foam.

Step C

The material from step B above (200 mg) in dry THF (5 mL) was treated at -40 °C under a nitrogen atmosphere with a 2 M solution of LDA in heptane/THF/benzene (Aldrich; 1.07 mL). Isobutyl iodide (0.09 mL) was then added, and the mixture left rise to room temperature. After 4 hr, the reaction mixture was quenched with saturated aqueous NH₄Cl and extracted with EtOAc. Washing with saturated aqueous NaHCO₃, drying over sodium sulfate and rotoevaporation afforded a mixture of the unreacted starting material and N-(tert-butoxycarbonyl)-3R-isobutyl-L-aspartic acid 4-piperidinamide.

Step D

The crude material from step C above in DMF (10 mL) was allowed to react with benzyl bromide (0.16 mL) and triethyl amine (0.24 mL) for 1 day at room temperature. After workup, the title compound was isolated by flash chromatography (gradient n-hexana/EtOAc mixtures) to obtain the title compound as a yellowish oil, which is the same as (4-benzyloxy-3S-tert-butoxycarbonylamino-2R-isobutyl)succinyl-piperidinamide obtained from the "azetidinone route" and described in Example 1 above. FT-IR (CHCl₃) 3433 br (NH), 1753 (ester CO), 1714 (carbamate CO), 1626 (amide CO) cm⁻¹. ¹H- NMR (200 MHz, CDCl₃) 0.89 and 0.91 (two d, 6 H, J= 6.6 Hz); 1.42 (s, 9 H); 1.2-1.8 (m, 9 H); 3.1-3.5 (m, 5 H); 4.48 (dd, 1 H, J= 4.1 and 10.0 Hz); 5.06 and 5.19 (two d, 2 H, J= 12.4 Hz); 6.45 (d, 1 H, J= 10.0 Hz); 7.32 (m, 5 H) ppm.

EXAMPLE 74

Tablets containing the following ingredients may be produced in a conventional manner:

	Ingredient	Per Tablet
30	Amine derivative	25.0 mg
	Lactose	125.0 mg
	Maize starch	75.0 mg

66

Talc	4.0 mg
Magnesium stearate	1.0 mg
Total weight	230.0 mg

EXAMPLE 75

- 5 Capsules containing the following ingredients may be produced in a conventional manner:


Ingredient	Per capsule
------------	-------------

Amine derivative	50.0 mg
Lactose	165.0 mg
Maize starch	20.0 mg
Talc	5.0 mg
Capsule fill weight	240.0 mg

10

CLAIMS

1. A compound which is an amine derivative of formula (I)



5

wherein

W is -CONHOH or -COOH;

R₁ and R₂, which are the same or different, are each hydrogen or

- a group G, which is methyl, C₂-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₃-C₇ cycloalkyl, cycloalkyl-C₁-C₁₀-alkyl, aryl, aryl-C₁-C₁₀-alkyl, aryl-C₂-C₁₀-alkenyl, heterocyclyl, heterocyclyl-C₁-C₁₀-alkyl or heterocyclyl-C₂-C₁₀-alkenyl, the said methyl, alkyl, alkenyl, cycloalkyl, aryl and heterocyclyl groups being unsubstituted or substituted by one to three substituents; or
- SO₂-G, wherein G is as defined above; or
- SO-G, wherein G is as defined above; or
- CO-G, wherein G is as defined above; or
- COO-G, wherein G is as defined above; or
- SO₂-NH₂, -SO₂-NHG or -SO₂-NGG', wherein G is as defined above and G', which is the same or different, is as defined above for G, or G and G', together with the nitrogen atom to which they are attached, form a saturated or unsaturated 3- to 7-membered azaheterocyclic ring, which may be fused to a carbocyclic, heterocyclic, or aromatic ring, and may be substituted at any carbon or additional nitrogen atom, or
- a group -CONH₂, -CONHG or -CO-NGG' wherein G and G' are as defined above, or G and G', together with the nitrogen atom to which they are attached, constitute a saturated or unsaturated 4- to 7-membered azaheterocyclic ring, which may be fused to a carbocyclic, heterocyclic, or aromatic ring, and may be substituted at any carbon or additional nitrogen atom, or

R₁ and R₂, taken together with the nitrogen atom to which they are attached, form a saturated or unsaturated 3- to 7-membered azaheterocyclic ring, which may be fused to a

carbocyclic, heterocyclic, or aromatic ring, and may be substituted at any carbon or additional nitrogen atom;

R₃ is C₁-C₁₅ alkyl, either unsubstituted or substituted by a C₃-C₇ cycloalkyl group, the alkyl and/or the cycloalkyl group being either unsubstituted, or substituted by one to three substituents selected from methyl, ethyl, C₃-C₄ linear or branched alkyl, fluoro, chloro, C₁-C₄ alkoxy, nitro, amino, dimethylamino, carboxy and carboxymethyl; or

R₃ is a group -R-X-R^I wherein R is a chemical bond, -CH₂-, -(CH₂)_m- wherein m is an integer from 2 to 5, -CH=CH-, -CH₂CH=CH-, phenylene (i.e., -C₆H₄-), -CH₂CH=CH-C₆H₄-, -CH₂CH₂CH=CH-, -CH₂-CC-, -CH₂CH₂-CC-, -CH₂CH₂CH=CH-C₆H₄-, -CH₂-CC-C₆H₄-, or -CH₂CH₂-CC-C₆H₄-; X is a direct bond, an oxygen atom, a sulfur atom, or a sulfinyl -S(O)-, sulfonyl -S(O)₂ or carbamoyl group -CONH- or -NHCO-; and R^I is C₁-C₆ alkyl, C₂-C₆ alkenyl, phenyl, phenyl-(C₁-C₆)-alkyl, phenyl-(C₂-C₆)-alkenyl, heterocyclyl, or heterocyclyl-(C₁-C₆)-alkyl, either unsubstituted or substituted by a group selected from F, Cl, Br, C₁-C₄ alkyl, C₁-C₄ alkoxy, alkylthio, arylthio, alkylsulfonyl, and arylsulfonyl;


Q, being a secondary or tertiary carboxamide, is:

- a group -CONHG or -CONGG', wherein G and G' are as defined above; or
 - a group -CONH-CHGG', wherein G and G' are as defined above; or a group -CONG''-CHGG', wherein G'', being the same or different, is defined as G above; or

- a group -CONH-CH₂-CHGG' or a group -CONG''-CH₂-CHGG', wherein G, G' and G'' are as defined above; or

- a group -CO-azaheterocyclyl, which is either unsubstituted or substituted at any carbon or additional nitrogen atom, with the proviso that when Q is -CONHG, and G is methyl, alkyl-methyl, cycloalkyl-methyl, aryl-methyl or heterocyclyl-methyl, then such methyl or substituted methyl cannot be further substituted by a group -(CH₂)_t-CO₂H, wherein t is 0, or esters and amides thereof; or a pharmaceutically acceptable salt, solvate or hydrate thereof.

2. A compound as claimed in claim 1 wherein the amine derivative has the formula (I'):



wherein:

W is -CONHOH or -COOH;

R₁ and R₂ are:

- both hydrogen; or

5 - both C₁-C₄ alkyl, or

R₁ is hydrogen or methyl, and R₂ is a group G which is:

- C₁-C₁₀ alkyl or C₂-C₁₀ alkenyl, either unsubstituted or substituted by C₃-C₇ cycloalkyl, or by one to three substituents selected from chloro, fluoro, hydroxy, C₁-C₄ alkoxy, C₁-C₄ alkyl, amino, methylamino, dimethylamino, -CONH₂, -CONHCH₃ or -CONHC(CH₃)₃, C₁-C₄ alkylthio, C₁-C₄ alkylsulfonyl, phenoxy, phenylthio and phenylsulfonyl, wherein the phenyl
10 group may in turn be substituted by chloro, fluoro, methoxy or methyl; or

- C₃-C₇ cycloalkyl; or

- an aryl group which is optionally substituted by one to three substituents selected from chloro, fluoro, hydroxy, C₁-C₄ alkoxy, amino, methylamino, dimethylamino, C₁-C₄ alkyl, C₃-
15 C₇ cycloalkyl, C₁-C₄ alkylthio, C₁-C₄ alkylsulfonyl, phenoxy, phenylthio and phenylsulfonyl, phenyl, benzyl, phenethyl, phenpropyl, naphthyl and pyridyl, and wherein any phenyl, naphthyl and pyridyl ring may in turn be substituted by one to three substituents selected from chloro, fluoro, methyl, hydroxy, methoxy, amino, methylamino and dimethylamino; or

- an unsaturated heterocyclyl group selected from pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, thiadiazolyl, thienyl, furyl, pyridyl, pyrazinyl, pyridazinyl, pyrimidinyl, benzothienyl, benzothiazolyl, benzoxazolyl, isobenzofuranyl, benzofuranyl, benzimidazolyl, indazolyl, chromenyl, indolyl, oxindolyl, quinolyl, isoquinolyl, isoindolyl, cinnolyl and purinyl, each of which is optionally substituted by one to three
20 substituents selected from chloro, fluoro, hydroxy, C₁-C₄ alkoxy, amino, methylamino, dimethylamino, C₁-C₄ alkyl, C₃-C₇ cycloalkyl, C₁-C₄ alkylthio, C₁-C₄ alkylsulfonyl, phenoxy, phenylthio and phenylsulfonyl, wherein the phenyl ring may be substituted by chloro, fluoro, methoxy or methyl; or

- a saturated or partially saturated heterocyclyl group selected from pyrrolidinyl, tetrahydrothienyl, tetrahydrofuryl, aziridinyl, oxiranyl, azetidyl, piperidinyl, piperazinyl, hexahydropyridazinyl, tetrahydropyranyl, 1-oxo-2-isoindolyl, tetrahydroisoquinolyl, hydantoinyl, morpholinyl, thiomorpholinyl, dioxanyl, dithianyl, azepinyl and thiazolidinyl, wherein any ring
30 nitrogen may be oxidized as an N-oxide, any ring carbon may be oxidized as a carbonyl, and any ring sulfur may be oxidized as a sulfoxide or sulfone; and the derivatives thereof substituted by one

to three substituents selected from chloro, fluoro, hydroxy, C₁-C₄ alkoxy, amino, methylamino, dimethylamino, C₁-C₄ alkyl, C₃-C₇ cycloalkyl, C₁-C₄ alkylthio, C₁-C₄ alkylsulfonyl, phenoxy, phenylthio and phenylsulfonyl, wherein the phenyl ring may be substituted by chloro, fluoro, methoxy or methyl; or

- 5 - C₁-C₁₀ alkyl, substituted by any of the unsaturated or saturated heterocyclyl groups as defined above, wherein any ring nitrogen may be oxidized as an N-oxide, any ring carbon may be oxidized as a carbonyl, and any ring sulfur may be oxidized as a sulfoxide or sulfone; or a derivative thereof substituted by one to three substituents selected from chloro, fluoro, hydroxy, C₁-C₄ alkoxy, amino, methylamino, dimethylamino, C₁-C₄ alkyl, C₃-C₇ cycloalkyl, -CONH₂, -CONHCH₃ and -CONHC(CH₃)₃, C₁-C₄ alkylthio, C₁-C₄ alkylsulfonyl, phenoxy, phenylthio and phenylsulfonyl, wherein the phenyl ring may be substituted by chloro, fluoro, methoxy or methyl; or

- 10 - C₁-C₁₀ alkyl, substituted by any of the aryl group as defined above, and the derivatives thereof substituted by one to three substituents selected from chloro, fluoro, hydroxy, C₁-C₄ alkoxy, amino, methylamino, dimethylamino, C₁-C₄ alkyl, C₃-C₇ cycloalkyl, C₁-C₄ alkylthio, C₁-C₄ alkylsulfonyl, phenoxy, phenylthio and phenylsulfonyl, wherein the phenyl ring may be substituted by chloro, fluoro, methoxy or methyl; or

R₁ is hydrogen or methyl, and R₂ is -SO₂-G, wherein G is as defined above; or

R₁ is hydrogen or methyl, and R₂ is -CO-G, wherein G is as defined above; or

- 20 R₁ is hydrogen or methyl, and R₂ is -CO-O-G, wherein G is as defined above; or

R₁ is hydrogen or methyl, and R₂ is -SO₂-NH₂, -SO₂-NHG or -SO₂-NGG', wherein G is as defined above and G', which is the same or different, is as defined above for G; or

R₁ is hydrogen or methyl, and R₂ is -SO-NH-G, wherein G is as defined above; or

- 25 R₁ is hydrogen or methyl, and R₂ is -CONHG or -CONGG', wherein G is as defined above and G', which is the same or different, is as defined above for G; or

R₁ and R₂, taken together with the nitrogen atom to which they are attached, form a 3- to 7-membered azaheterocyclyl ring, optionally containing N, O, S or SO₂ as an additional ring member, which may be substituted by oxo on one or two carbon ring atoms adjacent to the linking nitrogen atom, and which is optionally fused with a benzene ring, the azaheterocyclyl group being either unsubstituted or substituted at one or more carbon and/or nitrogen atoms by chloro, fluoro, hydroxy, C₁-C₄ alkoxy, amino, methylamino, dimethylamino, C₁-C₄ alkyl, phenyl, 4-fluorophenyl, benzyl, 4-fluorobenzyl, alpha-methylbenzyl, methylenedioxyphenyl, 2-phenethyl, 2-(4-fluorophenyl)ethyl, piperonyl, carbamoyl, -CONHCH₃, -CONHC(CH₃)₃, -

CONH-(4-fluorophenyl), -CONH-pyridyl, -CONH-(methylenedioxy)phenyl -CONH-piperonyl;
or

R₁ is hydrogen or methyl and R₂ is -SO₂-azaheterocyclyl, wherein azaheterocyclyl is as defined above; or

- 5 R₁ is hydrogen or methyl, and R₂ is -CO-azaheterocyclyl, wherein azaheterocyclyl is as defined above,


R₃ is -CH₂-alkyl, -(CH₂)_n-cycloalkyl, -(CH₂)_n-O-alkyl, -(CH₂)_n-O-(CH₂)_m-cycloalkyl, -(CH₂)_n-O-(CH₂)_m-aryl, -(CH₂)_n-O-(CH₂)_m-heterocyclyl, -(CH₂)_n-S-alkyl, -(CH₂)_n-S-(CH₂)_m-cycloalkyl, -(CH₂)_n-S-(CH₂)_m-aryl, -(CH₂)_n-S-(CH₂)_m-heterocyclyl, -(CH₂)_n-SO-alkyl, -
10 (CH₂)_n-SO-(CH₂)_m-cycloalkyl, -(CH₂)_n-SO-(CH₂)_m-aryl, -(CH₂)_n-SO-(CH₂)_m-heterocyclyl, -(CH₂)_n-SO₂-alkyl, -(CH₂)_n-SO₂-(CH₂)_m-cycloalkyl, -(CH₂)_n-SO₂-(CH₂)_m-aryl, -(CH₂)_n-SO₂-(CH₂)_m-heterocyclyl, -(CH₂)_n-CO-alkyl, -(CH₂)_n-CO-(CH₂)_m-cycloalkyl, -(CH₂)_n-CO-(CH₂)_m-aryl or -(CH₂)_n-CO-(CH₂)_m-heterocyclyl, wherein alkyl, cycloalkyl, aryl and heterocyclyl are as defined above, and n and m, being the same or different, are zero or an integer of 1 to 5,
15 and wherein the alkyl, cycloalkyl, aryl and heterocyclyl groups are optionally substituted by one to three substituents selected from chloro, fluoro, cyano, cyanomethyl, hydroxy, C₁-C₄ alkoxy, C₁-C₄ alkyl, C₁-C₄ alkylthio, C₁-C₄ alkylsulfonyl, phenyl, tolyl, 4-chlorophenyl, 4-fluorophenyl, 4-methoxyphenyl, 4-phenoxyphenyl, 4-(4-pyridyl)oxyphenyl, pyridyl; or R₃ is selected from isobutyl, n-hexyl, n-heptyl, n-octyl, n-nonyl, and cyclopentylmethyl; or R₃ is
20 selected from 2-phenylethyl, 3-phenylpropyl, 4-phenylbutyl, 5-phenylpentyl, wherein the phenyl group is either unsubstituted or substituted by chloro, fluoro, cyano, cyanomethyl, methyl, ethyl, propyl, butyl, mesyl, methoxy, ethoxy, propoxy, butoxy, phenoxy, 4-chlorophenoxy, 4-fluorophenoxy, benzyloxy, phenyl, 4-fluorophenyl, 4-chlorophenyl, 4-methoxyphenyl, 4-cyanophenyl, 4-cyanomethylphenyl, 4-pyridyl, 4-pyridyloxy; or R₃ is
25 selected from phenylsulfonylmethyl or phenylsulphonylethyl, wherein the phenyl group is either unsubstituted or substituted by chloro, fluoro, cyano, cyanomethyl, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, butoxy, phenoxy, 4-chlorophenoxy, 4-fluorophenoxy, benzyloxy, phenyl, 4-fluorophenyl, 4-chlorophenyl, 4-methoxyphenyl, 4-cyanophenyl, 4-cyanomethylphenyl, 4-pyridyl, 4-pyridyloxy;

- 30 Q is:

- a group -CONHG, -CONGG', -CONH-CHGG', CON(CH₃)-CHGG', -CONH-CH₂-CHGG', or -CON(CH₃)-CH₂-CHGG', wherein G and G', being as defined above, are selected from C₁-C₆ straight or branched alkyl, C₃-C₆ cycloalkyl, phenyl, tolyl,

- methylenedioxyphenyl, piperonyl and pyridyl, either unsubstituted or substituted by one to three substituents selected from chloro, fluoro, hydroxy, hydroxymethyl, C₁-C₄ alkoxy, amino, methylamino, dimethylamino, C₁-C₄ alkyl, C₃-C₇ cycloalkyl, -CONH₂, -CONHCH₃, -CONHC(CH₃)₃, -CONH(4-fluorophenyl), -CONH-pyridyl, -CONH-(methylenedioxy)phenyl,
- 5 -CONH-piperonyl, carbomethoxy, carbethoxy, or a keto group -CO-R^{II}, wherein R^{II}, being as defined above, is selected from C₁-C₄ alkyl, phenyl, fluorophenyl, chlorophenyl, methylenedioxyphenyl, naphthyl, piperonyl, or a sulfone -(CH₂)_n-SO₂-R^{II}, wherein n and R^{II} are as defined above, or a sulfonamide -(CH₂)_n-SO₂-NH₂, -(CH₂)_n-SO₂-NHR^{II}, -(CH₂)_n-SO₂-NR^{II} R^{III}, wherein n, R^{II} and R^{III} are as defined above, including the special case wherein R^{II} and R^{III},
- 10 and R^{III}, taken together with the nitrogen atom to which they are attached, constitute an azaheterocyclyl ring, as defined above; or
- a group -CO-azaheterocyclyl, wherein azaheterocyclyl, being as defined above, is either unsubstituted or substituted by one to three substituents selected from hydroxy, hydroxymethyl, C₁-C₄ alkoxy, carbamoyl, carbomethoxy, carbethoxy, mesyl, C₁-C₆ linear or
- 15 branched alkyl, trifluoromethyl, C₃-C₇ cycloalkyl, aryl, heterocyclyl and aryl-(C₁-C₃)alkyl or heterocyclyl-(C₁-C₃)alkyl; or said azaheterocyclyl group is substituted by a group -CONH-R^{II}, wherein R^{II}, being as defined above, is selected from methyl, isopropyl, tert-butyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, 4-fluorophenyl, 4-chlorophenyl, 4-methoxyphenyl, 3,4-difluorophenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 4-piperidyl, 2-thiazolyl, 1-
- 20 naphthyl, 2-naphthyl, 3-quinolyl, 5-quinolyl, 3-isoquinolyl, 5-isoquinolyl, 3-quinuclidinyl, methylenedioxyphenyl, piperonyl, 2-benzimidazolyl and 5-tetrazolyl; or said azaheterocyclyl group is substituted by a keto group -CO-R^{II}, or by a carbinol group of formula -CH(OH)-R^{II}, wherein R^{II}, being as defined above, is selected from C₁-C₄ alkyl, phenyl, fluorophenyl, chlorophenyl, methylenedioxyphenyl, naphthyl, piperonyl, or by a sulfone -(CH₂)_n-SO₂-R^{II},
- 25 wherein n and R^{II} are as defined above, or by a sulfonamide -(CH₂)_n-SO₂-NH₂, -(CH₂)_n-SO₂-NHR^{II}, -(CH₂)_n-SO₂-NR^{II} R^{III}, wherein n, R^{II} and R^{III} are as defined above, including the special case wherein R^{II} and R^{III}, taken together with the nitrogen atom to which they are attached, constitute an azaheterocyclyl ring, as defined above;
- and the salts and solvates thereof.
- 30 3. A compound of formula (II'), which is a cyclic prodrug of a compound of formula (I') as claimed in claim 2:


73



wherein R_2 , R_3 and Q are as defined in claim 2, T is methyl or a hydrogen atom, and T' is methyl, C_2 - C_4 lower alkyl, phenyl, benzyl, optionally substituted by one to three substituents
 5 selected from C_1 - C_4 alkyl, hydroxy, C_1 - C_4 alkoxy, amino, methylamino, dimethylamino, chloro and fluoro.

4. A compound as claimed in claim 2 wherein R_3 is selected from isobutyl, n-hexyl, n-heptyl, n-octyl, n-nonyl, cyclopentylmethyl; or is selected from 3-phenylpropyl, 4-phenylbutyl, 5-phenylpentyl, 3-(p-diphenyl)propyl, 3-(4-pyridyl-4-phenyl)propyl, phenoxymethyl, 2-phenoxyethyl, 2-(phenylmethoxy)ethyl, 3-phenoxypropyl, 4-(phenylmethoxy)butyl, phenylsulfonylmethyl, 2-(phenylsulfonyl)ethyl, 3-(4-benzoylphenyl)propyl, including
 10 derivatives thereof substituted at any phenyl carbon atom by one to three substituents selected from chloro, fluoro, hydroxy, C_1 - C_4 alkyl and C_1 - C_4 alkoxy.


5. A process for producing a compound as defined in claim 1, comprising:
 15 (a) reacting a beta-lactam compound of formula (III):




wherein R_2 and R_3 are as defined in claim 1 and W' is $COOH$, $CONHOH$, or a protected derivative thereof, with a primary or secondary acyclic amine of formula $G-NH_2$, $GG'NH$,
 20 $GG'CH-NH_2$, $GG'CH-NHCH_3$, $GG'CH-CH_2-NH_2$, or $GG'CH-CH_2-NHCH_3$, wherein G and G' are as defined in claim 1, or with an amine of formula $G-NH_2$, $GG'NH$, $GG'CH-NH_2$ or $GG'CH-CH_2-NH_2$, wherein G and G' are as defined above; including the special case

74


wherein G and G', taken together with the nitrogen atom to which they are attached, constitute an azaheterocyclyl ring, wherein azaheterocyclyl is as defined in claim 1, to obtain a compound of formula (IV):



- 5 wherein W', R₂ and R₃ are as defined above and Q is defined as in claim 1; and
 (b) converting said compound of formula (IV) into a compound of formula (I):




- 10 wherein W, R₁, R₂, R₃ and Q are as defined in claim 1; and
 (c) if desired, removing the protecting groups and/or if desired, converting any of the groups W, R, R₁, R₂, R₃ and Q into different groups W, R, R₁, R₂, R₃ and Q at the end or at any stage of the process.
6. A process for producing a compound as defined in claim 2, which process comprises
- 15 (a) reacting a beta-lactam compound of general formula (III'):



- wherein R₂ and R₃ are as defined in claim 2, and W' is COOH or CONHOH, or a protected derivative thereof, with an amine as defined in claim 5, to obtain a compound of formula
- 20

75


(IV'):



wherein W', R₂, R₃ and Q are as defined in claim 2; and

(b) converting this compound of formula (IV') into a compound of formula (I'):

5




wherein W, R₁, R₂, R₃ and Q are as defined in claim 2; and

(c) if needed, removing the protecting groups and/or, if desired, converting any of the groups W, R, R₁, R₂, R₃ and Q into different groups W, R, R₁, R₂, R₃ and Q at the end or at any stage of the process.

7. A process for producing a compound of formula (I) as defined in claim 1, which process comprises:

(a) reacting an aspartic acid derivative of formula (V):


15



wherein W' is either carboxy or a protected derivative thereof, Q' is either Q as described above, or carboxy or protected carboxy, and R₁, R₂ are as defined in claim 1, with a reagent of formula R₃-X, wherein R₃ is as defined in claim 1, and X is chloro, bromo or iodo, or sulfonyloxy, e.g. triflate or mesylate,


20 to obtain a compound of formula (VI):

76



wherein W', R₁, R₂, R₃ and Q' are as described above; and

- 5 (b) converting of such compound of formula (VI) into a compound of formula (I):



wherein W, R₁, R₂, R₃ and Q are as described above; and

- 10 (c) if needed, removing the protecting groups and/or, if desired, converting any of the groups W, R, R₁, R₂, R₃ and Q into different groups W, R, R₁, R₂, R₃ and Q at the end or at any stage of the process.
8. A pharmaceutical composition which comprises a pharmaceutically acceptable diluent or carrier and, as an active ingredient, a compound as defined in any of claims 1 to 4.
- 15 9. A compound as defined in any one of claims 1 to 4 for use in a method of treatment of the human or animal body by therapy or prophylaxis..
10. A compound as claimed in claim 9 for use as an inhibitor of a matrix metalloproteinase.
11. A compound as claimed in claim 9 for use as an inhibitor of the release of TNF- α from cells.
- 20 12. A compound as claimed in any one of claims 9, 10 and 11 for use in the therapeutic or prophylactic treatment of a tumoural or inflammatory disease.

INTERNATIONAL SEARCH REPORT

Int'l Application No

PCT/EP 98/04220

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D295/18 A61K31/445 C07C259/06 A61K31/16 A61K31/19
C07C237/04 C07C237/06 C07D233/32 C07D317/68 C07D213/74
C07D405/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07C C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 108, no. 13, 28 March 1988 Columbus, Ohio, US; abstract no. 112899, BURGER, KLAUS ET AL: "A new route to 3-alkyl-substituted aspartic acid derivatives" XP002083194 see compounds of formula V & CHEM.-ZTG. (1986), 110(11), 422-4, 1986, ---	1,2
A	WO 96 33166 A (DU PONT MERCK PHARMA) 24 October 1996 see claim 1; example 3; table 1 --- -/--	1,2,8-12

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

2 December 1998

Date of mailing of the international search report

28/12/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

De Jong, B

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/04220

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO 95 33731 A (HOFFMANN LA ROCHE ;BROADHURST MICHAEL JOHN (GB); BROWN PAUL ANTHON) 14 December 1995 see claim 1</p> <p>-----</p>	1,2,8-12

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/04220

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9633166 A	24-10-1996	US 5691381 A	25-11-1997
		AU 5556396 A	07-11-1996
		CA 2218380 A	24-10-1996
		EP 0821669 A	04-02-1997
WO 9533731 A	14-12-1995	AU 2615695 A	04-01-1996